



Europäisches Patentamt
European Patent Office
Office européen des brevets

Publication number:

**0 292 977
A1**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 88108449.5

(22) Date of filing: 27.05.88

(51) Int. Cl.: C07D 311/22 , C07D 311/24 ,
C07D 311/58 , C07D 311/66 ,
A61K 31/35 , C07C 57/48 ,
C07C 61/39

(30) Priority: 29.05.87 US 57136
12.05.88 US 188708

(43) Date of publication of application:
30.11.88 Bulletin 88/48

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI NL SE


(71) Applicant: G.D. Searle & Co.
P.O. Box 5110
Chicago Illinois 60680(US)

(72) Inventor: Djuric, Stevan Wakefield
1023 Huber Lane
Glenview, IL 60025(US)
Inventor: Shone, Robert Larry
1441 Joan Drive
Palatine, IL 60067(US)
Inventor: Yu, Stella Siu Tzyy
7801 Maple Street
Morton Grove, IL 60053(US)

(74) Representative: Wolff, Hans Joachim, Dr.jur.
Dipl.-Chem. et al
Bell, Wolff & Bell Rechtsanwälte Postfach 80
01 40 Adelonstrasse 58
D-6230 Frankfurt am Main 80(DE)

(54) Alkoxy-substituted dihydrobenzopyran-2-carboxylate derivatives.

(57) This invention encompasses compounds of Formula 1 and the pharmaceutically acceptable salts thereof.



wherein

R¹ represents alkyl having 2-6 carbon atoms;

R² represents methyl or ethyl;

R³ represents alkyl having 1 to 5 carbon atoms;

EP 0 292 977 A1

W represents $(CH_2)_x$ where x is 2 to 7, alkylene having 2 to 7 carbon atoms, alkenylene having 3 to 7 carbon atoms, alkynylene having 3 to 7 carbon atoms, or cyclopentyl;

R⁴ represents hydrogen, alkyl having 2-5 carbon atoms, alkenyl having 2 to 5 carbon atoms, or alkynyl having 2 to 5 carbon atoms;

Q represents oxygen or CH_2 ;

B represents CH_2 , $C=O$ or $CH-OH$;

R⁵ represents hydrogen, alkyl having 1 to 6 carbon atoms, or R⁵ and R⁶ together optionally represent a carbon to carbon bond; or R⁵ represents alkanoyl having 2 to 4 carbon atoms, carboxy, alkoxy, carbonyl, or $(CH_2)_y-CO_2R^8$ wherein y is 0 to 4 and R⁸ is hydrogen or alkyl having 1 to 6 carbon atoms; and

A represents $-Z-CO_2R^7$ or $-Z-COR^9R^{10}$ wherein R⁷ represents hydrogen or alkyl having 1 to 6 carbon atoms, R⁹, R¹⁰, represent hydrogen, alkyl having 1 to 6 carbon atoms, or cycloalkyl having 3 to 6 carbon atoms or NR^9R^{10} form a heterocyclic ring, and * in Z is absent or represents straight or branched chain alkylene or alkenylene having up to 6 carbon atoms.

These compounds are selective antagonists of leukotriene B₄ (LTB₄) with little or no antagonism of leukotriene D₄ (LTD₄) and are useful anti-inflammatory agents for treating inflammatory bowel disease, rheumatoid arthritis, gout and psoriasis.

ALKOXY-SUBSTITUTED DIHYDROBENZOPYRAN-2-CARBOXYLATE DERIVATIVES

BACKGROUND OF THE INVENTION

5 This is a continuation-in-part of United States Serial No. 07/057,136 filed May 29, 1987.

Field of the Invention

10 This invention is in the field of pharmaceutical agents which selectively act as leukotriene B₄ (LTB₄) antagonists.

Prior Art

15 The prior art discloses compounds having structures similar to Formula I except in the prior art compounds R² of Formula I is replaced with hydrogen. For example:

EPA 079,637
 EPA 100,466,6 (U.S.4,665,203)
 20 U. S. Patent 4,595,882
 U. S. Patent 4,546,194
 EPA (U.S.S.N.) 507383 129,906
 CA 103 (19) 160 389 0
 Japan 60/42378
 25 EPA 150447

Journal of Medicinal Chemistry, 1977, 20 (3) 376 broadly discloses compounds where R² of Formula I is hydrogen.

EPA 79,647 generically discloses a formula which encompasses compounds of Formula I wherein -OR² is -O-alkyl and B is C=O, but does not exemplify or otherwise enable the preparation and use of such
 30 compounds. EPA 79,637 extensively discloses intermediates to making compounds of this invention, that is, where -OR² in formula I is -OH. EPA 79,637 does not teach the selective LTB₄ antagonist activity of compounds of this invention.

U.S. 4,281,008, U.S. 3,822,148, and U.S. 4,006,245 generically disclose formulae which encompass compounds of Formula I wherein -OR² is -O-alkyl or O-Me and B is C=O but do not exemplify or otherwise
 35 enable the preparation and use of such compounds, nor do they teach the selective LTB₄ antagonist activity of compounds of the present invention.

The Journal of Medicinal Chemistry, 1977, Vol. 20 (3): 376 discloses a compound similar to the compounds of Formula I except the acyl and alkyl substituents adjacent the -OR₂ in Formula I are absent. The Journal of Medicinal Chemistry article also discloses a compound similar to compounds of Formula I
 40 except for a hydroxy substituent on the -O-(CH₂)_x-O- connecting group.

The prior art generally describes the above compounds as LTD₄ antagonists for use as anti-allergy compounds or as antagonists of SRS-A, the slow reacting substance of anaphylaxis. In sharp contrast compounds of Formula I are selective LTB₄ antagonists useful in treating inflammatory diseases.

Leukotriene D₄ and C₄ (LTD₄/LTC₄) and leukotriene B₄ (LTB₄) are products of the arachidonic acid
 45 metabolic pathway. LTD₄ and LTC₄ are associated with smooth muscle contraction and contract guinea pig ileum, human and guinea pig bronchi and human pulmonary artery and vein. LTB₄ is associated with neutrophil stimulation and is characterized by chemotaxis, aggregation and degranulation. LTB₄ is believed to be an important mediator of inflammation. High levels of LTB₄ are detected in rheumatoid arthritis, gout, psoriasis, and inflammatory bowel disease. Thus antagonists of LTB₄ are useful in the therapy of such
 50 diseases.

Gastroenterology, 1985: 88 : 580-7 discusses the role of arachidonic acid metabolites in inflammatory bowel disease.


British Medical Bulletin, (1983) vol. 39, No. 3, pp. 249-254, generally discusses the pharmacology and

pathophysiology of leukotriene B₄.

Biochemical and Biophysical Research Communications, Vol. 138, No. 2 (1986), pp. 540-546 discusses the pharmacology of a specific LTB₄ antagonist which has a different structure than compounds of this invention.

BRIEF DESCRIPTION OF THE INVENTION

This invention encompasses compounds of Formula I and the stereoisomers and pharmaceutically acceptable salts thereof;



I

wherein

R¹ represents alkyl having 2-6 carbon atoms, alkenyl having 2 to 6 carbon atoms, alkynyl having 2 to 6 carbon atoms, or -(CH₂)_n-R wherein R represents cycloalkyl of 3 to 5 carbon atoms and n is 1 or 2;

R² represents methyl or ethyl;

R³ represents alkyl having 1 to 5 carbon atoms;

W represents (CH₂)_x where x is 2 to 7, alkenylene having 3 to 7 carbon atoms, alkynylene having 3 to 7 carbon atoms, or cyclopentyl;

R⁴ represents hydrogen, alkyl having 2 to 5 carbon atoms, alkenyl having 2 to 5 carbon atoms, or alkynyl having 2 to 5 carbon atoms;

Q represents oxygen or CH₂;

B represents CH₂, C=O, or CH-OH;

R⁵ represents hydrogen, alkyl having 1 to 6 carbon atoms, or R⁵ and R⁶ together optionally represent a carbon to carbon bond; or R⁵ represents alkanoyl having 2 to 4 carbon atoms, carboxy, alkoxycarbonyl, or (CH₂)_y-CO₂R⁸ wherein y is 0 to 4 and R⁸ is hydrogen or alkyl having 1 to 6 carbon atoms; and


A represents -Z-CO₂R⁷ or -Z-CONR⁹R¹⁰ wherein R⁷ represents hydrogen or alkyl having 1 to 6 carbon atoms, R⁹ and R¹⁰ represent hydrogen, alkyl having 1 to 6 carbon atoms, or cycloalkyl having 3 to 6 carbon atoms, or NR⁹R¹⁰ form a heterocyclic ring, and wherein Z is absent or represents straight or branched chain alkylene or alkenylene having up to 6 carbon atoms.

These compounds are selective antagonists of leukotriene B₄ (LTB₄) with little or no antagonism of leukotriene D₄ (LTD₄) and are useful anti-inflammatory agents for treating inflammatory bowel disease, rheumatoid arthritis, gout and psoriasis.

DETAILED DESCRIPTION OF THE INVENTION

This invention encompasses the compounds of formula I as previously described.

Preferred embodiments of the present invention are compounds of the formula Ia, the stereoisomers and pharmaceutically acceptable salts thereof,



Ia

wherein

R¹ represents alkyl having 2-4 carbon atoms;

R² represents methyl or ethyl;

R³ represents alkyl having 1 to 3 carbon atoms;

x represents 3 to 5;

R⁴ represents alkyl having 2 to 4 carbon atoms;


B represents CH₂ or C=O;

R⁵ represents hydrogen, alkyl having 1 to 4 carbon atoms, or R₅ and R₆ together optionally represent a carbon to carbon bond;

A represents -Z-CO₂R⁷, wherein R⁷ represents hydrogen or alkyl having 1 to 4 carbon atoms and wherein Z is absent or represents alkylene having up to 2 carbon atoms.

These compounds are selective antagonists of leukotriene B₄ (LTB₄) with little or no antagonism of leukotriene D₄ (LTD₄) and are useful anti-inflammatory agents for treating inflammatory bowel disease, rheumatoid arthritis, gout, and psoriasis.


More preferred embodiments are compounds of the formula II



II


where R₅, A, B, R₆ are as previously defined in formula Ia.

Most preferred embodiments of the present invention are compounds of the formula IIa, the stereoisomers and pharmaceutically acceptable salts thereof



wherein p is 0 to 2.

A particularly preferred embodiment is the compound of the formula III:



III

7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid

and the pharmaceutically active salts thereof. This compound and its pharmaceutically acceptable salts are particularly preferred because of the specificity of the LTB₄ antagonist activity and its oral bioavailability.

Other compounds encompassed by this invention are as follows:

7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-4-oxo-8-propyl-2H-1-benzopyran-2-propanoic acid

7-[[5-(4-acetyl-3-methoxy-2-propylphenoxy)pentyl]oxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid, ethyl ester

7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoic acid

7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-propanoic acid

7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-2-carboxy-3,4-dihydro-4-oxo-8-propyl-2H-1-benzopyran-2-propanoic acid

7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-2-(methoxycarbonyl)-4-oxo-8-propyl-2H-1-benzopyran-2-propanoic acid, methyl ester

7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-2-methyl-8-propyl-2H-1-benzopyran-2-propanoic acid

7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-2-methyl-8-propyl-2H-1-benzopyran-2-propanoic acid, methyl ester

7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoic acid

7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoic acid, methyl ester

7-[3-(4-acetyl-2-(cyclopropylmethyl)-3-methoxyphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid

5 7-[[5-(4-acetyl-3-methoxy-2-propylphenoxy)pentyl]oxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid

Alkyl defined for R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹ and R¹⁰ is straight or branched chain alkyl having the indicated number of carbon atoms.

Pharmaceutically acceptable salts such as ammonium, sodium, potassium, alkaline earth, tetraalkylammonium and the like are encompassed by the invention.

Compounds of this invention are generally prepared by alkylating the prior art phenol hydroxy (R² = H) compounds to form compounds of formula I by conventional techniques. Thus, the reaction of the phenol hydroxy (R² = H) with methyl iodide in potassium carbonate provides the ether. Dimethyl sulfate in acetone and base is also useful in preparing ethers. Alternatively intermediates can be alkylated prior to forming the -O-(CH₂)-O- bridge. Hydrolysis of the ester compounds in the presence of lithium hydroxide and methanol gives the acid compounds.

The biological activity of compounds of this invention is indicated by the following tests.

20 Preparation of Human Neutrophils

Neutrophils were purified from venous blood of normal human donors using standard techniques of dextran sedimentation, centrifugation on Ficoll-paque® (Pharmacia) or Histopaque® sterile solution (Sigma) and hypotonic lysis of erythrocytes (Boyum, A., Isolation of Leukocytes From Human Blood: Further
 25 Observations, Scand. J. Lab. Clin. Invest., 21 (Suppl. 97): 31, 1968). The purity of isolated neutrophils was >95%.

LTB₄ Receptor Binding Assay

30 Neutrophils (4 - 6x10⁶) in 1ml Hanks' balanced salt solution containing 10 mM HEPES buffer (HBSS), pH 7.4 and 30μm nordihydroguaiaretic acid were incubated with 0.6x10⁻⁹ M (³H) LTB₄ in the presence or absence of test compounds. The incubation was carried out at 0°C for 45 minutes and terminated by adding 5ml of ice-cold HBSS followed by rapid filtration of incubation mixture under vacuum through GF/C
 35 glass fiber filters. The filters were further washed with 10ml HBSS and radioactivity was determined. Specific binding was defined as the difference between total binding and nonspecific binding which was not displaced by 10⁻⁷ M unlabeled LTB₄. All data refer to specific binding.

40 Human Neutrophil Degranulation Assay

Neutrophil degranulation was determined by measuring the release of myeloperoxidase activity into the incubation medium. Neutrophils (3 x 10⁶) in 1 ml HBSS solution were preincubated with cytochalasin B (5 μg) at 37°C for 5 minutes, followed by preincubation with test compounds for 7 minutes. Neutrophils were
 45 then incubated for 2 to 20 minutes with either LTB₄ (5 x 10⁻⁸ M) or the chemotactic peptide f-met-leu-phe (5 x 10⁻⁶ M) to induce degranulation. Following incubation, samples were centrifuged and myeloperoxidase was extracted from the cell pellets by sonication in phosphate buffer containing 0.4% Triton X-100. Triton X-100 was also added to the supernatants to a concentration of 0.4%. The supernatants and the pellet - extracts were then assayed spectrophotometrically for myeloperoxidase activity by determining the rate of
 50 decomposition of H₂O₂ with o-dianisidine as hydrogen donor as described by Renlund, et al. (Renlund, D. G., MacFarlane, J. L., Christensen, R. D., Lynch, R. E., and Rothstein, G., A Quantitative And Sensitive Method For Measurement of Myeloperoxidase, Clinical Research 28:75A, 1980), Myeloperoxidase activity released into the supernatant was expressed as the percent of the average total activity (pellet plus supernatant).

55

Guinea Pig LTB₄-Induced Dermal Chemotaxis

Test compound was administered intravenously or intragastrically at various times prior to the injection of leukotriene B₄ (LTB₄). LTB₄ was diluted in phosphate buffered saline (PBS) and 35 mg in 0.2ml were injected intradermally into the shaven backs of anesthetized guinea pigs. PBS was injected as control. Four hours later, animals were sacrificed, skins removed and stored frozen (-70 °C). Injection sites were removed with a skin punch and mechanically homogenized (Polytron, Brinkmann Instruments). Myeloperoxidase (MPO), a marker enzyme for neutrophils, was extracted with 0.5% hexadecyltrimethylammonium bromide in 50 mM potassium phosphate buffer (pH 6.0), using sonication and freeze-thaw procedures. After centrifugation (40,000 x g, 30 minutes), enzyme activities in the supernatants were assayed spectrophotometrically (A₄₆₀) by measuring the decomposition of hydrogen peroxide with o-dianisidine after 15 minutes. MPO activity was found to be proportional to the number of neutrophils. In guinea pigs the level of MPO activity increased with the amount of LTB₄ injected.

Modified Boyden Chamber Chemotaxis

Human neutrophils were isolated from citrated peripheral blood using standard techniques of dextran sedimentation, followed by centrifugation on Histopaque® sterile solution (Sigma) or Ficoll-paque® (Pharmacia) and hypotonic lysis of erythrocytes. A final cell suspension of 3.4×10^6 neutrophils/ml of HEPES-buffered Hanks' balanced salt solution (HBSS, pH 7.3) was added to the upper well (0.8ml) of a modified Boyden chamber (blind well). The lower well (0.2ml), separated by a polycarbonate membrane (Nucleopore Corp.), contained HBSS or 3×10^{-8} M LTB₄ in the presence or absence of test compound. Following a 90 minute incubation at 37 °C in 5% CO₂-95% air, cells from the lower well were lysed and nuclei counted in a Model S-Plus-IV Coulter Counter. Percent inhibition was calculated from cell counts corrected for random migration by subtracting the mean of the HBSS control.

The compounds of this invention can be administered in a number of dosage forms. A preferred method of delivery would be oral or in such a manner so as to localize the action of the inhibitor. In an inflammatory condition such as rheumatoid arthritis the compounds could be injected directly into the affected joint. The compounds could also be administered in oral unit dosage forms such as tablets, capsules, pills, powders or granules. They may be introduced intraperitoneally, subcutaneously, or intramuscularly using forms known to the pharmaceutical art. Topical application in the form of salves and ointments are useful for treating psoriasis. Regardless of the route of administration selected, the compounds are formulated into pharmaceutically acceptable dosage forms by conventional methods known to the pharmaceutical art.

LTD₄ Antagonism In Guinea Pig Ileum

Fresh segments of guinea pig ileum were suspended in 2ml. tissue baths containing oxygenated modified Tyrodes solution. After an equilibration period, an agonist dose-response curve was generated by exposing each tissue to 4 different LTD₄ doses and measuring the ensuing contractile heights. The ileum segments were washed and rested between exposures to agonist. Following this, the tissues were incubated with a single concentration of test compound and the agonist dose-response procedure was repeated. The dose ratio is a measure of the antagonist's ability to shift the agonist dose-response curve to the right. It is derived as the concentration of agonist necessary to reach a given response level in the presence (A') versus the absence (A) of antagonist. For example, if the test concentration of compound had no effect on the agonist-induced response (A' = A) the dose-ratio would approximate 1. Dose-ratios increase if the compound inhibits the agonist-induced response. One dose-ratio value is determined for each strip of ileum used to test antagonist. If the dose-ratios increase as a function of increasing antagonist concentration, these data may be evaluated by Schild analysis to determine whether the inhibition is competitive and if so, what the pA₂ value is for the compound. Schild analysis examines the linearity of the function described by the dose-ratios written as log (A'/A) versus antagonist concentration. If linearity is confirmed and the slope approximates -1, inhibition is considered to be competitive. The pA₂ is the negative log of the antagonist concentration required to produce a dose-ratio of 2. This value is considered to be a measure of the affinity of the competitive antagonist.

Leukotriene D₄ Receptor Assay Using Guinea Pig Lung Preparations

The following description discloses an LTD₄ radioreceptor assay used to identify compounds which inhibit LTD₄ binding to lung membrane preparations.

MEMBRANE PREPARATIONS:

For each membrane preparation, 10-11 male Hartley guinea pigs, weighing less than 350 gm, were sacrificed by guillotine. The lungs were rapidly removed and placed in ice-cold 50 mM Trisma 7.0 buffer. After all the lungs were collected, each was probed free of bronchial tissue to at least the level of tertiary branching as observed through a dissecting lamp. The tissue was minced with a razor blade on a damp paper towel and placed into fresh ice-cold Trisma 7.0 buffer. When all the tissues had been processed, they were pooled, drained and weighed. Nine volumes of cold buffer were added and the tissue was polytroned for 6-10 second bursts with a 1 minute cool down period in an ice bath between each burst. The homogenate was filtered through a gauze pad and centrifuged at 1085 x g for 15 minutes at 4 °C. Supernatants were saved and centrifuged at 40,000 x g for 20 minutes at 4 °C. This time, supernatants were discarded and the pellets were resuspended in 30-35 ml fresh Trisma 7.0 buffer by 1-10 second bursts with the polytron. The materials were centrifuged again at 40,000 x g. Supernatants were again discarded. Each pellet was resuspended again with approximately 7 ml buffer. All this material was pooled into a 100 ml polypropylene container and stirred with a magnetic stirring bar. The tubes were rinsed with small amounts of buffer and the rinse was added to the membrane preparation. The homogenized membranes were aliquoted into 1.5 ml microfuge tubes and frozen at -70 °C. Three aliquots were submitted for protein determinations by FPB (fluorescamine protein binding) assay. This procedure usually resulted in protein concentrations ranging from 1-5 mg/ml.

RADIORECEPTOR ASSAY METHODS:

Each guinea pig lung preparation yielded enough protein to last through several months of assaying. Because of this, each preparation was tested for its 3H-LTD₄ dissociation constant, number of receptor populations, number of binding sites per unit protein, and the dose-response displacement characteristics by LTD₄ and the receptor antagonist FPL 55712.

Unless otherwise indicated, all binding experiments were performed in a final incubation volume of 250 µl. Tritiated LTD₄ was obtained from New England Nuclear and nonradioactive LTD₄ was purchased from Biomol Research Laboratories, Inc. (Philadelphia, PA). Both materials were received in solutions. 3H-LTD₄ was solubilized in 50% EtOH with 0.01M pH 6.8 phosphate buffer while Biomol LTD₄ was prepared in 65% MeOH with 35% water and small amounts of AcOH and NH₄OH. Both solutions were brought to appropriate assay concentrations by dilution in assay buffer consisting of 50 mM Trisma 7.4 with 5 mM L-cysteine and 20 mM CaCl₂. Membranes were also diluted with this assay buffer. Compounds other than LTD₄ were solubilized in DMSO and added to each tube in 5 µl aliquots. Diluent was added to all other appropriate control tubes. Incubations were conducted for 30 minutes at 25 °C in a shaking water bath. Termination of the reaction was achieved by pouring the incubate over 2.5 cm Whatman GF/C filters soaked in assay buffer and set over the ports of Millipore filter vacuum manifolds. Filters were rinsed with 4 ml ice cold assay buffer 3 times. The filters were removed, placed in 10 ml Aquasol scintillation cocktail and allowed to cool in the dark for 2 hours prior to counting. All samples were corrected for background and isotope decay before converting DPM determination to mass.

DETERMINATION OF IC₅₀ VALUES:

LTD₄ specific binding was determined as the difference between 3H-LTD₄ binding alone and in the presence of 1 mM unlabeled LTD₄. Nonspecific LTD₄ binding was also removed from tubes containing compound as well. The percent inhibition of binding for each dose of compound was calculated by subtracting specific binding in the presence of compound from LTD₄ specific binding in the absence of compound. This result was then divided by the specific binding without compound and multiplied by 100. These values were logit-transformed and linear regression was performed on the dose-response data. All

dose-response sets used to calculate IC_{50} values contained data from 3-6 compound concentrations. Correlation coefficients for the straight line model almost always exceeded 0.98. IC_{50} values were calculated from the regression equation for the line.

The compounds of the present invention are formulated into pharmaceutically acceptable dosage forms by conventional methods known to those skilled in the art. The compounds may be administered in a number of dosage forms, for example, such oral dosage forms as tablets, capsules, pills, powders, or granules. They may also be administered intravascularly, intraperitoneally, subcutaneously, or intramuscularly using forms known to the pharmaceutical art.


An effective but non-toxic quantity of the compound is employed in treatment. The dosage regimen for inhibition of LTB_4 by the compounds of this invention is selected in accordance with a variety of factors including the type, age, weight, sex, and medical condition of the mammal, the particular disease and its severity, the route of administration and the particular compound employed. An ordinarily skilled physician or veterinarian will readily determine and prescribe the effective amount of the compound to prevent or arrest the progress of the condition. In so proceeding, the physician or veterinarian could employ or use relatively low dosages at first, subsequently increasing the dose until a maximum response is obtained. A dosage range of 1 to 25 mg/kg generally provides a therapeutically effective anti-inflammatory response.

The following examples illustrate the preparation of compounds of this invention from known starting materials. The invention, which is set forth in the foregoing disclosure, is not to be construed or limited either in spirit or in scope by these examples. Those skilled in the art will readily understand that known variations of the conditions and processes of the following preparative procedures can be used to prepare these compounds. All temperatures are degrees Celsius unless otherwise noted.

U.S. 4,665,203 issued May 12, 1987 discloses methods for making some of the intermediates used in making compounds of the present invention.


Example 1

7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.



III

(a) 493mg of methyl 7-[3-(4-acetyl-3-hydroxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate was added to 25ml of acetone containing 276 mg of anhydrous potassium carbonate and 282mg of methyl iodide. The mixture was refluxed for about 24 hours and water was added and the mixture was then extracted with ethyl acetate. The extract was dried, the solvent removed under vacuum, and the residual oil was chromatographed over silica gel with a 40/60 mixture of ethyl acetate/hexane to provide pure methyl ether, methyl 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate having the formula



1 a

(b) The methyl ether (1a) (340 mg) was dissolved in methanol (5ml) containing lithium hydroxide (0.7ml of a 2N LiOH solution in water). The mixture was stirred at room temperature overnight and the solvent removed in vacuo. The residue was partitioned between ethyl acetate and 2N HCl and the organic layer separated and washed with brine. Evaporation of the volatiles in vacuo afforded crude acid of Formula III. This material was purified by silica gel chromatography using ethyl acetate/hexane/acetic acid (40:60:0.5) as eluant. The pure product was recrystallized from ethyl acetate/hexane to afford 200 mg of product, 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid, m.p. 65-68° C.

Microanalysis:

Found

C 69.22, H 7.53

Theory

C 69.40, H 7.49

The NMR (CDCl₃) shows a -OCH₃ at δ 3.75.

Example 2Comparative Test Data For Compound III

5

10

15

20

25

30

35

40

45

50

	<u>LTB₄ Receptor</u>	<u>Guinea Pig Ileum</u>
	<u>Binding</u>	<u>Contraction pA₂</u>
Compound III	50% at 0.3 μ M	6.5 or less
Compound III where -OCH ₃ is replaced with -OH	30% at 1 μ M	7.6


Guinea Pig Ileum Results

<u>Compound</u>	<u>Test</u> <u>Concentration (M)</u>	<u>Dose</u> <u>Ratio</u>
Compound III where -OCH ₃ is replaced with -OH	1x10 ⁻⁶	100
	3x10 ⁻⁷	13.24
		9.97
		17.04
		10.59
	1x10 ⁻⁷	3.48
		7.38
	3x10 ⁻⁸	1.71
		1.97
		3.10
Compound III		1.89
	3x10 ⁻⁶	8.65
		11.16
	1x10 ⁻⁶	2.54
		1.95
	3x10 ⁻⁷	2.28
		1.67


Thus, the -OCH₃ compound (III) is about 5x more potent than the -OH compound in LTB₄ receptor binding and at least 10x less active as an LTD₄ antagonist in the guinea pig ileum smooth muscle contraction test. Compounds of this invention are selective LTB₄ antagonists useful in treating inflammatory disease.

Example 3

7-[3-[4-acetyl-2-(cyclopropylmethyl)-3-methoxyphenoxy]propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.




(a) A mixture of Zinc/copper couple (509 mg, 7.8mMol), iodine (2 mg), and methylene iodide (2.09g, 7.8 mMol) was suspended in ether (5ml) and the mixture heated to reflux for 30 minutes. The heat was removed and 2,4-dihydroxy-3-allylacetophenone (500 mg, 2.6mMol) was added as a solution in ether. The mixture was heated at reflux for 2 hours and a further 2g of methylene iodide was added during this time. The mixture was cooled and stirred at room temperature for 10 hours. The mixture was filtered and the ether solution washed sequentially with aqueous ammonium chloride, sodium bicarbonate and brine. The organic layer was dried over sodium sulfate and evaporated in vacuo to afford a crude residue which was purified by column chromatography on silica gel (eluting with 2:8 ethyl acetate/hexane to give 220 mg of 2,4-dihydroxy-3-cyclopropylmethylacetophenone
NMR ^1H (δ , CDCl_3) 0.25-0.5 (4H, m, cyclopropyl H's), 1.0 (1H, m, cyclopropyl H), 2.5 (3H, s, acetyl CH_3), 2.65(2H, d, CH_2 - ϕ) 6.4(1H, d, ArH), 7.5(1H, d, ArH)



3a


(b) 1,3 chlorobromopropane (1.9g), methyl 7-hydroxy-8-propylchroman-2-carboxylate (2.5g) and potassium carbonate (2.1g) were stirred at RT in dry dimethylformamide under argon for 24 hours. The mixture was poured into water and thoroughly extracted with ethyl acetate. The combined organic extracts were washed sequentially with water and brine and then dried over magnesium sulfate. Filtration and evaporation of the volatiles in vacuo afforded 4.3g of crude product which was chromatographed on silica gel (10:90 ethyl acetate/hexane) to provide 3.0g of methyl 7-(3-chloropropoxy)-8-propylchroman-2-carboxylate.
NMR ^1H (δ , CDCl_3) 0.8(3H, t, propyl CH_3) 1.5-2.6 (8H, m propyl CH_2 's and ring CH_2 's), 3.7 (3H, s, CO_2CH_3), 3.75 (2H, t, CH_2Cl), 4.2 (2H, t, CH_2OAc), 4.75(1H, d, CH-OAr) 6.4 (1H, d, ArH), 6.8 (1H, d, ArH)



3b


(c) 2,4-dihydroxy-3-cyclopropylmethylacetophenone (0.2g), methyl 7-(3-chloropropoxy)-8-propylchroman-2-carboxylate (327mgs), sodium iodide (0.15g) and potassium carbonate (0.14g) were stirred in 3 ml dry dimethylformamide (DMF) overnight at 45° under argon. The mixture was poured into water and thoroughly extracted with ethyl acetate. The combined organic extracts were washed with water and dried over magnesium sulfate. Evaporation of the volatiles in vacuo afforded a crude oil which was purified by chromatography on silica gel to give 98 mgs of methyl 7-[3-[4-acetyl-2-(cyclopropylmethyl)-3-hydroxyphenoxy]propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate.

NMR: ^1H (δ , CDCl_3) 0.2-0.45 (4H, m, cyclopropyl H's), 0.9 (3H, t, propyl CH_3), 1.0 (1H, m, cyclopropyl H), 1.4-2.8 (12H, m, aliphatic and cyclic CH_2 's), 2.55 (3H, s, CH_3CO), 3.75 (3H, s, CO_2CH_3), 4.25 (2H, t, CH_2OAr), 4.35 (2H, t, CH_2OAr), 4.75 (1H, m, CH-OAr), 6.4-7.6 (4H, m, ArH 's)



3c

(d) Compound 3c (95 mg) was dissolved in acetone (5ml) containing dimethyl sulfate (38 mg) and potassium hydroxide (12.4 mg) and the mixture was heated at 56° C under argon for 10 hours. The solvent was removed in vacuo and the residue partitioned between ethyl acetate and water. The organic layer was separated, washed with potassium carbonate solution and dried over magnesium sulfate. Evaporation of the volatiles in vacuo afforded a crude oil which was purified by chromatography on silica gel (1:9 ethyl acetate/hexane) to give 40 mg of pure methyl 7-[3-[4-acetyl-2-(cyclopropylmethyl)-3-methoxyphenoxy]propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate. NMR ^1H (δ , CDCl_3) 3.5 (3H, s, OCH_3).



3d

(e) Compound 3d (38 mg), was dissolved in 0.3ml of methanol containing 74 μl of a 1M lithium hydroxide solution, and the solution was stirred overnight at room temperature. The solvent was removed

and the residue partitioned between ethyl acetate and 10% hydrochloric acid solution. The organic layer was removed and dried over magnesium sulfate. Evaporation of the volatiles in vacuo afforded 35 mg of the title compound, 7-[3-[4-acetyl-2-(cyclopropylmethyl)-3-methoxyphenoxy]propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.

5 Micronanalysis:

Found:

C 69.51, H 7.34.

Calculated for $C_{25}H_{36}O_7 \cdot 1/4H_2O$:

C 69.30, H 7.49.

10


Example 4

15

7-[[3-(4-acetyl-3-methoxy-2-propylphenoxy)cyclopentyl]oxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.

20

25




30

(a) 2,4-dihydroxy-3-propyl acetophenone (1.94g), 1,3-dihydroxycyclopentane (1.02g), triphenylphosphine (2.62g), and diethyl azodicarboxylate (1.7g) were dissolved in dry tetrahydrofuran (THF) (200 ml) and stirred under argon at room temperature overnight. The solvent was evaporated and the crude residue purified by silica gel chromatography (Merck 60, 4:6 ethyl acetate/hexane) to afford 1.5g of 1-(2-propyl-3-hydroxy-4-acetylphenoxy)-3-hydroxycyclopentane.

35

NMR 1H (δ , $CDCl_3$) 0.9 (3H,t,propyl CH_3), 1.3-2.6(10H, aliphatic and cyclic CH_2 's), 2.55 (3H,s, CH_3CO), 4.4 (1H, m, CH-OH), 4.9 (1H, m, CH-OAr), 6.4 (1H, d, ArH), 7.6 (1H, d, ArH)

40




45

(b) Compound 4a (1.4 mg, 5.0 mMol), methyl 7-hydroxy-8-propylchroman-2-carboxylate (1.33 mg, 5.01mMol), triphenylphosphine (1.32 mg, 5.0 mMol) and diethyl azodicarboxylate (875 mg) were dissolved in dry tetrahydrofuran (120 ml) and stirred under argon at room temperature for 10 hours. The solvent was evaporated and the solid residue dissolved in dry ether and filtered through sintered glass. The filtrate was evaporated and the residue purified by chromatography on silica gel using 2:8 ethyl acetate/hexane as eluant. 0.7 g of methyl 7-[[3-(4-acetyl-3-hydroxy-2-propylphenoxy)cyclopentyl]oxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate were obtained.

55


NMR 1H (δ , $CDCl_3$) 0.9(6H,t, propyl CH_3 's), 1.5-2.7 (18H, aliphatic and cyclic CH_2 's), 2.55(3H,s, CH_3CO), 3.75 (3H,s, OCH_3), 4.75(1H,m, $\underline{CH-CO_2CH_3}$), 4.9(1H,m, CH-OAr), 5.0(1H, m,CH-OAr), 6.4 (2H, 2ds, ArH's) 6.8- (1H,d,ArH), 7.55(1H,d,ArH)



4b

(c) Compound 4b (419 mg, 0.8 mMol) was dissolved in acetone (10ml) containing potassium hydroxide (55.3 mg, 0.98mMol), and dimethyl sulfate (134.5 mg). The mixture was heated at 45 °C for 10 hours and then cooled and the volatiles removed in vacuo. The residue was partitioned between ethyl acetate and water and the organic layer separated and dried over magnesium sulfate. Removal of solvent afforded a crude oil which was purified using radial band chromatography (Harrison chromatotron, 2:8 ethyl acetate/hexane as eluant). Methyl 7-[[3-(4-acetyl-3-methoxy-2-propylphenoxy)cyclopentyl]oxy]-3,4-dihydro-

8-propyl-2H-1-benzopyran-2-carboxylate was obtained pure as a colorless oil (270 mg). NMR ^1H (δ , CDCl_3) 0.9(6H, 2t's, propyl CH_3 's), 1.5-2.8 (18H, aliphatic and cyclic CH_2 's), 2.6 (3H, s, CH_3CO), 3.75 (3H, s, OCH_3 ether), 3.77 (3H, s, OCH_3 ester), 4.75 (1H, m, CHCO_2CH_3), 4.9 (1H, m, CH-OAr), 5.0 (1H, m, CH-OAr), 6.4(1H, d, ArH), 6.65 (1H, d, ArH), 6.8(1H, d, ArH), 7.55 (1H, d, ArH)




4c

(d) Compound 4c (150 mg) was dissolved in methanol (2ml) containing 1M lithium hydroxide (450 μL) and the mixture was stirred at room temperature overnight. The volatiles were removed in vacuo and the residue partitioned between ethyl acetate and dilute hydrochloric acid. The organic layer was separated and dried over magnesium sulfate. Removal of solvent afforded the title compound, 7-[[3-(4-acetyl-3-methoxy-2-propylphenoxy)cyclopentyl]oxy]-3,4-dihydro-3-propyl-2H-1-benzopyran-2-carboxylic acid. NMR ^1H (δ , CDCl_3) no OCH_3 ester at 3.77.


Example 5

7-[3-[4-acetyl-3-methoxy-2-(2-propenyl)phenoxy]propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.



(a) 2,4-Dihydroxy-3-(2-propenyl) acetophenone (Aldrich, 12.5g, 65mMol) and methyl 7-(3-iodopropoxy)-8-propylchroman-2-carboxylate (29.9g, 71.5mMol) were dissolved in DMF (300ml) containing powdered potassium carbonate (26.91g, 195mMol). The mixture was allowed to stir at room temperature under nitrogen for 8 hours and then partitioned between water and toluene. The organic layer was separated and washed with water and brine and then dried over sodium sulfate. Evaporation of the volatiles in vacuo afforded 36g of crude product which was chromatographed on Merck 60 silica gel using 5:95 ethyl acetate/toluene as eluant. 25g of pure methyl 7-[3-[4-acetyl-3-hydroxy-2-(2-propenyl)phenoxy]propoxy]-3,4-


dihydro-8-propyl-2H-1-benzopyran-2-carboxylate were obtained in this manner.
NMR ^1H (δ , CDCl_3) 0.9(3H,t,propyl CH_3), 1.4-2.8(OH, aliphatic and cyclic CH_2 's), 2.55 (3H,d, CH_3CO), 3.45- (2H,d, $\text{ArCH}_2\text{CH}=\text{CH}_2$), 3.75 (3H,s, CO_2CH_3), 4.09(2H,t, CH_2OAr), 4.22(2H,t, CH_2OAr) 4.75(1H,t, CHOAr), 4.85-5.00(2H,m,allyl H's), 5.75-6.1(1H,m,allyl CH) 6.4-7.12(4H,ArH's)



5a

(b) Compound 5a (20.6g, 42.7mMol) was dissolved in dry DMF (105ml) containing methyl iodide (18.4g, 128.5mMol) and powdered potassium carbonate (17.69g, 128mMol). The mixture was stirred under nitrogen at room temperature for 22 hours and then partitioned between water and toluene. The organic layer was separated, washed with water and brine and dried over sodium sulfate. Removal of the volatiles in vacuo afforded 21g of product which was methyl 7-[3-[4-acetyl-3-methoxy-2-(2-propenyl)phenoxy]propoxy]-3,4-

dihydro-8-propyl-2H-1-benzopyran-2-carboxylate.
NMR ^1H (δ , CDCl_3) 3.77 (3H, s, OCH_3 ether)



5b

(c) Compound 5b (16.96g, 34.15mMol) was dissolved in methanol and a solution of lithium hydroxide in water (4.3g in 43ml 102.5mMol) was added. The mixture was stirred at room temperature for 3 hours and the solvent removed in vacuo. The mixture was partitioned between toluene and dilute hydrochloric acid and the organic layer removed. The organic extract was washed sequentially with water, brine and then dried over sodium sulfate. Evaporation of the volatiles in vacuo afforded a yellow oil which was purified by silica gel chromatography on Biosil A using 20% ethyl acetate in toluene as eluant. The product, 7-[3-[4-acetyl-3-methoxy-2-(2-propenyl)phenoxy]propoxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid was isolated as a light yellow oil, 15g, which was recrystallized from ethyl acetate/hexane, mp 85.5° C.

Microanalysis:

Found,


C 69.69, H 7.10.

Calculated for C₂₈H₃₅O₇,


C 69.6, H 6.98.

Example 6

7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid




(a) 2,4-dihydroxyacetophenone (7.1g, 0.05 mol) and dimethyl oxylate (7.2g) were dissolved in DMF containing a solution of sodium methoxide in methanol (4.0g Na, 100ml MeOH) and the mixture was stirred at room temperature for 48 hours. At this point, acetic acid (180ml) was added and the mixture was heated to 100° C for 5 hours. The solvent was removed in vacuo and the residue partitioned between ethyl acetate and water. The organic layer was separated, the aqueous layer was thoroughly extracted with more ethyl acetate, and the combined organic extracts were washed with brine. Evaporation of the dried (Na₂SO₄) solvent in vacuo afforded methyl 7-hydroxy-4-oxo-4H-1-benzopyran-2-carboxylate (6a) as a crude yellow solid which was crystallized from ethyl acetate/hexane to afford 3.5g of pure product. NMR: ¹H (δ, CDCl₃) 4.0 (3H, s, CO₂CH₃), 6.85 (1H, s, chromenone H) 6.9 - 8.0 (3H, Ar H's)



6a


(b) Compound 6a (3.57g, 16mMol) was dissolved in ethyl acetate containing phosphoric acid and 5% palladium on carbon. The mixture was shaken in a Parr apparatus at room temperature under an atmosphere of hydrogen for 22 hours. The solution was filtered and washed with water and the organic layer was dried over magnesium sulfate. Evaporation of the volatiles in vacuo afforded a crude oil which was purified by silica gel chromatography using ethyl acetate/hexane 6:4 as eluant. 3.3g of product, methyl 7-hydroxy-chroman-2-carboxylate, (6b) was obtained as a clear oil. NMR: ¹H (δ, CDCl₃) 2.1-2.8 (4H, m, cyclic CH₂'s) 3.8(3H,s, CO₂CH₃), 4.7 (1H, dd, CHOAr), 6.35-6.9(3H, Ar H's)



6b

(c) Compound 6b (416 mg, 2mMol) and 3-(2-propyl-3-hydroxy-4-acetylphenoxy)-1-iodopropane (720mg, 2mMol) were dissolved in dry DMF (3ml) containing powdered anhydrous potassium carbonate (552 mg, 4mMol) and the mixture was stirred at 60 °C for 10 hours. The mixture was cooled and partitioned between ethyl acetate and water. The organic layer was separated, washed with water and dried over magnesium sulfate. Evaporation of the volatiles in vacuo afforded a crude oil which was purified by flash chromatography on Merck 60 silica gel using ethyl acetate and hexane (2.5:7.5) as eluant. 540 mg of product, methyl 7-[3-(4-acetyl-3-hydroxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate, (6c) were obtained.


NMR ¹H (δ, CDCl₃) 0.9 (3H, t, propyl H's), 1.4-2.8 (10H, cyclic CH₂'s and aliphatic CH₂'s), 2.55(3H,s,CH₃CO) 3.79 (3H,s,OCH₃), 4.2 (2H,t,OA), 4.2 (2H,t,OA), 4.7 (1H,m,CH-OAr), 6.4-7.6(5H,ArH's)



6c

(d) Compound 6c (250mg) was dissolved in distilled acetone (5ml) containing potassium hydroxide (41.8 mg) and dimethyl sulfate (117.6 mg). The mixture was heated with stirring under argon at 40 °C for 10 hours. The mixture was cooled, the solvent removed and the residue partitioned between ethyl acetate and water. The organic layer was separated and dried over magnesium sulfate. Evaporation of the volatiles in vacuo afforded 240 mg of product, methyl 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate, (6d) which was homogeneous by thin layer chromatography.

NMR ¹H(δ, CDCl₃) 3.65 (3H, s, OCH₃)



(e) Compound 6d (200 mg) was dissolved in methanol (3ml) containing lithium hydroxide (1.0 ml of a 1M solution) and the mixture was stirred at room temperature for 10 hours. The solvent was removed and the residue partitioned between ethyl acetate and dilute hydrochloric acid. The organic layer was separated, washed with brine and dried over sodium sulfate. Evaporation of the volatiles in vacuo afforded 40 mg of

product, 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-4-oxo-8-propyl-4H-1-benzopyran-2-carboxylic acid, as a gum.

Microanalysis:

Found


5 C 67.59, H 6.93.

Calculated for $C_{25}H_{30}O_7$,

C 67.86, H 6.83.


Example 7

15 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-4-oxo-8-propyl-4H-1-benzopyran-2-carboxylic acid



30 (a) Methyl 7-[3-(4-acetyl-3-hydroxy-2-propylphenoxy)propoxy]-4-oxo-8-propyl-4H-1-benzopyran-2-carboxylate (250mg, 0.48mMol) was dissolved in acetone (10ml) containing potassium hydroxide (35 mg, 0.53mMol), and dimethyl sulfate (78.3 mg 0.62mMol). The mixture was heated at reflux temperature for 10 hours and then cooled. The residue was partitioned between ethyl acetate and water and the organic layer separated and dried over sodium sulfate. The solvent was removed in vacuo to afford 260 mg of product, methyl 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-4-oxo-8-propyl-4H-1-benzopyran-2-carboxylate.

35 NMR: $^1H(\delta: CDCl_3)$ 0.9(6H, 2t's, propyl CH_3 's), 1.4-2.8 (10H, aliphatic CH_2 's), 3.75(3H,s, OCH_3 ether), 4.0-(3H,s, CH_3 ester), 4.25(2H,t, CH_2OAr), 4.35 (2H,t, CH_2OAr) 7.05(1H,s, olefinic H) 6.7, 7.05, 7.55 and 8.05 (4H, d's, Ar H's)



7a

50 (b) Compound 7a (200 mg, 0.4mMol) was dissolved in methanol (2ml) containing 0.48ml of a 1M lithium hydroxide solution. The mixture was stirred at room temperature for 10 hours and the solvent removed. The residue was partitioned between ethyl acetate and dilute hydrochloric acid and the organic layer was separated, dried over magnesium sulfate, filtered and evaporated in vacuo. The product, 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-4-oxo-8-propyl-4H-1-benzopyran-2-carboxylic acid, was isolated as a light yellow gum (122 mg, 62% yield).

Microanalysis:

Found


C 66.43, H 6.59.

Calculated for $C_{28}H_{32}O_8 \cdot 1/2H_2O$;

C 66.43, H 6.59.

Example 8

7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-N-methyl-8-propyl-2H-1-benzopyran-2-carboxamide



The compound of Example 1 (80 mg,) was treated with a slight excess of oxalyl chloride and the mixture left to stand at room temperature for 2 hours. The mixture was stripped, dissolved in methylene chloride and methylamine gas was bubbled into the reaction mixture. The mixture was washed with water, dried over magnesium sulfate filtered and the solvent was evaporated in vacuo. The resultant white solid was washed with ethyl acetate/hexane (2:8) and dried in vacuo. 60 mg of product was obtained.

Microanalysis:

Found


C 69.63, H 7.96, N 2.86,

Calculated for $C_{29}H_{35}NO_6$,

C 70.00, H 7.9, N 2.82.

Example 9

Methyl 7-[[5-(4-acetyl-3-methoxy-2-propylphenoxy)pentyl]oxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate.



Methyl 7-[[5-(4-acetyl-3-hydroxy-2-propylphenoxy)pentyl]oxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate (1mMol), prepared as described in U.S. 4,565,882, was dissolved in acetone containing 2.5 equivalents of potassium hydroxide and 3 equivalents of dimethyl sulfate. The mixture was heated at 40 °C

for 10 hours and then cooled; the solvent was removed in vacuo and the residue partitioned between ethyl acetate and water. The organic layer was separated and dried over magnesium sulfate. Evaporation of the volatiles in vacuo afforded the methyl ether.


NMR ^1H (δ , CDCl_3)

- 5 0.9 (6H, 2t's, propyl CH_3 's), 1.4-2.8 (18H, aliphatic and cyclic CH_2 's), 2.55 (3H, s, CH_3CO), 3.75 (3H, s, OCH_3) 3.85 (3H, s, OCH_3), 3.9 (2H, t, CH_2OAr), 4.1 (2H, t, CH_2OAr), 4.7 (1H, dd CHOAr) 6.4, 6.65, 6.8, and 7.55 (4H, ArH's)

The compounds of Examples 10 to 14 were prepared according to the procedure described in Example 9 beginning with the appropriate phenol.

Example 10

Methyl 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-l-benzopyran-2-propanoate.




The title compound was prepared according to the method of Example 9 from methyl 7-[3-(4-acetyl-3-hydroxy-2-propylphenoxy)propoxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-l-benzopyran-2-propanoate, which was prepared as described in U.S. 4,665,203.

NMR ^1H (δ , CDCl_3)

- 35 0.9 (6H, 2ts, propyl CH_3 's), 1.4-2.7 (16H, aliphatic and cyclic CH_2 's), 1.35 (3H, s, CH_3), 2.6 (3H, s, CH_3CO), 3.7- (3H, s, OCH_3), 3.75 (3H, s, OCH_3), 4.25 (4H, CH_2OAr), 6.6, 6.7, 7.6 and 7.75 (4H, d's, ArH's)

Example 11

Methyl 7-[[4-(4-acetyl-3-methoxy-2-propylphenoxy)-2-butynyl]oxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-l-benzopyran-2-propanoate.




The title compound was prepared by the method of Example 9 from methyl 7-[[4-(4-acetyl-3-hydroxy-2-propylphenoxy)-2-butynyl]oxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-l-benzopyran-2-propanoate.

NMR ^1H (δ , CDCl_3)

0.9 (6H, 2t's, propyl CH_3 's), 1.4-2.7 (14H, aliphatic and cyclic CH_2 's), 1.35 (3H, s, CH_3), 2.6 (3H, s, CH_3CO), 3.7 (3H, s, OCH_3), 3.75 (3H, s, OCH_3), 4.8 (4H, s, CH_2OAr), 6.6, 6.7, 7.65 and 7.75 (4H, d's, ArH's)

Example 12

Methyl 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-1,2,3,4-tetrahydro-8-propyl-2-naphthalenecarboxylate.



(a) 1.6 grams of 3-(2-n-propyl-3-hydroxy-4-acetylphenoxy)-1-bromopropane (.005 mole) which was prepared as described in U.S. Patent 4,565,882 (Miyano et al.) was dissolved in 30 ml of methyl ethyl ketone. Two grams (2.5 equivalents) of potassium carbonate was added to the reaction along with 1.25 grams (.005 mole) of methyl 1,2,3,4-tetrahydro-7-hydroxy-8-propyl-2-naphthalenecarboxylate obtained in Example 28. After 100 mg of sodium iodide was added, a nitrogen blanket was placed on the system and the contents were refluxed for forty-eight hours. The reaction was cooled to room temperature. The mixture was filtered and the filtrate evaporated. The residue was chromatographed on silica gel with elution by 3% acetone-toluene. Eluant evaporation left 1.45 grams (60%) of compound (12a), methyl 7-[3-(4-acetyl-3-hydroxy-2-propylphenoxy)propoxy]-1,2,3,4-tetrahydro-8-propyl-2-naphthalenecarboxylate, m.p. $84-85^\circ\text{C}$.


Microanalysis:

Calc:

C 72.17, H 7.94; for $\text{C}_{29}\text{H}_{38}\text{O}_6$

Found:

C 71.87, H 7.90.



12a


(b) The title compound was prepared by the method of Example 9 from methyl 7-[3-(4-acetyl-3-hydroxy-2-propylphenoxy)propoxy]-1,2,3,4-tetrahydro-8-propyl-2-naphthalenecarboxylate.

NMR ^1H (δ , CDCl_3)

0.95(6H, 2t's, propyl CH_3 's), 1.4-3.1 (16H, aliphatic and cyclic CH_2 's), 2.6 (3H, s, CH_3CO), 3.75 (6H, 2 overlapping singlets, OCH_3 's), 4.5 (2H, t, CH_2OAr), 4.25 (2H, t, CH_2OAr), 6.7, 6.9, 7.55 (4H, Ar H's)

Example 13

Methyl 3,4-dihydro-7-[3-[3-methoxy-4-(2-methyl-1-oxopropyl)-2-propylphenoxy]propoxy]-8-propyl-2H-1-benzopyran-2-carboxylate.



(a) A solution of 495 mg (1.89 mMole) of triphenylphosphine, 420 mg (1.89 mMole) of 2,4-dihydroxy-3-propylisobutyrophenone, 582 mg (1.89 mMole) of methyl 7-(3-hydroxypropoxy)-8-n-propylchroman-2-carboxylate in 5 ml of tetrahydrofuran was prepared and cooled in an ice bath. Diethyl azocarcboxylate, 350 mg (1.89 mMole) was added and the solution was allowed to warm to room temperature and stir for 18 hours.

The solvent was removed by rotary evaporation and the residue triturated with 25 ml ether, cooled and filtered. The ether filtrate was then washed with water, brine, and dried over anhydrous magnesium sulfate. The drying agent was removed by filtration, the ether removed by rotary evaporation, and the residue purified by elution chromatography on silica gel with 1% acetone-toluene to furnish 720 mg (75%) of the desired ester, methyl 3,4-dihydro-7-[3-[3-hydroxy-4-(2-methyl-1-oxopropyl)-2-propylphenoxy]propoxy]-8-propyl-2H-1-benzopyran-2-carboxylate, as an oil.


Microanalysis:

Calc:

C, 70.08; H, 7.86;

Found:


C, 69.51; H, 7.81.



(b) The title compound was prepared by the method of Example 9 from methyl 3,4-dihydro-7-[3-(3-hydroxy-4-(2-methyl-1-oxopropyl)-2-propylphenoxy)propoxy]-8-propyl-2H-1-benzopyran-2-carboxylate.
 NMR ^1H (δ , CDCl_3)
 0.9(6H, 2t's, propyl CH_3 's), 1.15 (6H, d, $(\text{CH}_3)_2\text{CCO}$), 1.4-2.7 (14H, aliphatic and cyclic CH_2 's), 3.7 (3H, s, OCH_3) 3.75 (3H, s, OCH_3), 4.15-4.2 (4H, 2t's, CH_2OAr), 4.75 (1H, 1dd, CHOAr), 6.4, 6.7, 6.8 and 7.35 (4H, d's, ArH's)

Example 14

Ethyl 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-propanoate.




The title compound was prepared by the method of Example 9 from ethyl 7-[3-(4-acetyl-3-hydroxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-propanoate which was prepared as described in U.S. 4,665,203.

NMR ^1H (δ , CDCl_3)
 0.9 (6H, 2ts, propyl CH_3 's), 1.2-2.7 (20H, aliphatic cyclic and $\text{CO}_2\text{CH}_2\text{CH}_3$ H's), 2.6 (3H, s, CH_3CO), 3.75 (3H, s, OCH_3), 4.0 (1H, m, CHOAr), 4.1, 4.3 (4H, 2 triplets, CH_2OAr), 6.4, 6.7, 6.8, 7.55 (4H, ArH's)

Example 15


7-[[5-(4-acetyl-3-methoxy-2-propylphenoxy)pentyl]oxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.



The compound prepared in Example 9 (1mMol) was dissolved in methanol containing two equivalents of a 1M lithium hydroxide solution in water. The mixture was stirred at room temperature for 10 hours and then solvent was removed in vacuo. The residue was partitioned between ethyl acetate and water, the organic layer was separated and dried over magnesium sulfate and the volatiles were removed in vacuo. The product carboxylic acid may be purified by column chromatography on silica gel if necessary. NMR ^1H (δ , CDCl_3) 0.9 (6H, 2ts, propyl CH_3 's), 1.4-2.8 (18H, aliphatic and cyclic CH_2 's), 2.55 (3H, s, CH_3CO), 3.75 (3H, s, OCH_3), 3.9 (2H, t, CH_2OAr), 4.1 (2H, t, CH_2OAr) 4.75 (1H, m, CHOAr), 6.4, 6.65, 6.8, and 7.55 (4H, ArH's)

Example 16

7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoic acid.



The title compound was prepared from the compound of Example 10 using the method of Example 15.

Microanalysis:

Found:


C 68.55, H 7.60.

Calculated for $\text{C}_{31}\text{H}_{40}\text{O}_8$:

C 68.87, H 7.46.

Example 17


7-[[4-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-2-butyryloxy-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoic acid.



The title compound was prepared from the compound of Example 11 using the method of Example 15. NMR ^1H (δ , CDCl_3) 0.9 (6H, 2t's, propyl CH_3 's), 1.4-2.7 (14H, aliphatic and cyclic CH_2 's), 1.35 (3H, s, CH_3), 2.6 (3H, s, CH_3CO), 3.75 (3H, s, OCH_3), 4.85 (4H, brs, CH_2OAr), 6.6, 6.7, 7.6, and 7.75 (4H, d's, ArH's)

Example 18


7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-1,2,3,4-tetrahydro-8-propyl-2-naphthalenecarboxylic acid.



The title compound was prepared from the compound of Example 12 using the method of Example 15. Microanalysis:
Found: C 71.85, H 7.94.
Calculated for $\text{C}_{24}\text{H}_{28}\text{O}_5$:
C 72.17, H 7.94.

Example 19


3,4-dihydro-7-[3-[3-methoxy-4-(2-methyl-1-oxopropyl)-2-propylphenoxy]propoxy]-8-propyl-2H-1-benzopyran-2-carboxylic acid.



The title compound was prepared from the compound of Example 13 using the method of Example 15.
NMR ^1H (δ , CDCl_3) 0.9 (6H, 2t's, propyl CH_3 's), 1.15 (6H, d, $(\text{CH}_3)_2\text{CCO}$), 1.4-2.7 (14H, aliphatic and cyclic CH_2 's), 3.7 (3H, s, OCH_3), 4.15, 4.2 (4H, 2t's, CH_2OAr) 4.75 (1H, dd, CHOAr), 6.4, 6.7, 6.82 and 7.35 (4H, d's, ArH 's)

Example 20


7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-propanoic acid.




The title compound was prepared from the compound of Example 14 using the method of Example 15.
NMR ^1H (δ , CDCl_3) 0.9 (6H, 2t's, propyl CH_3 's), 1.2-2.7 (18H, aliphatic and cyclic CH_2 's), 2.6 (3H, s, CH_3CO), 3.75 (3H, s, OCH_3), 4.0 (1H, m, CHOAr), 4.1, 4.3 (4H, 2t's, CH_2OAr), 6.4, 6.7, 6.8, 7.55 (4H, ArH 's)

Example 21

7-[3-(4-acetyl-3-ethoxy-2-propylphenoxy)-propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.




(a) 4-(3-chloropropoxy)-2-ethoxy-3-propylacetophenone (5.0g, 18.47mMol), potassium carbonate (7.3g), and iodoethane (3.0ml) were dissolved in dry DMF (50ml) and the mixture stirred overnight at room temperature. The reaction mixture was then partitioned between ether and water and the organic layer separated. The ether extracts were washed with water and brine and then dried over sodium sulfate. Evaporation of the volatiles in vacuo afforded 5.36g of crude 4-(3-chloropropoxy)-2-ethoxy-3-propylacetophenone.




21a

(b) 4-(3-chloropropoxy)-2-ethoxy-3-propylacetophenone (5.36g) was dissolved in acetone containing 10 equivalents of sodium iodide, and the mixture was refluxed under nitrogen for 10 hours. Solvent was removed in vacuo and the residue partitioned between ether and water. The organic extracts were washed with water and brine, dried over sodium sulfate and evaporated in vacuo to afford 7.0g of a yellow oil, 4-(3-iodopropoxy)-2-ethoxy-3-propylacetophenone. NMR ^1H (δ , CDCl_3) 300 MHz 3.4 (2H, t, CH_2I)



21b

(c) Compound 21b (2.34g, 6mMol), methyl 7-hydroxy-8-propylchroman-2-carboxylate (1.25g, 5mMol) and potassium carbonate (2.07g, 15mMol) were suspended in dry DMF (12.5ml) and the mixture stirred at room temperature for 48 hours. The mixture was then partitioned between ether and water and the ether layer separated, washed with water and brine and then dried over sodium sulfate. Evaporation of the volatiles in vacuo afforded 2.9g of crude product which was purified by chromatography on silica gel to afford 1.34g of pure methyl 7-[3-(4-acetyl-3-ethoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate.



21c

(d) Compound 21c (0.8g, 1.56mMoles) was dissolved in methanol/water (7:3, 25ml) containing lithium hydroxide (0.13g, 3.12mMol). The mixture was stirred at room temperature for 2 hours and then partitioned between dilute hydrochloric acid and ether. The organic layer was separated, washed with brine, dried over sodium sulfate and concentrated in vacuo to afford 1.1g of crude product. This material was purified by chromatography on silica gel using ethyl acetate/hexane/acetic acid (50:50:1) as eluant. 7-[3-(4-acetyl-3-ethoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2- carboxylic acid was obtained as 0.71g of white crystals m.p. 81-84 ° C.

Microanalysis:

Found


C 69.68, H 7.85.

Calculated for $C_{30}H_{40}O_7$,

C 69.85, H 7.68.

Example 22

7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-4-oxo-8-propyl-4H-1-benzopyran-2-propanoic acid



Starting with methyl 7-[3-(4-acetyl-3-hydroxy-2-propylphenoxy)propoxy]-4-oxo-8-propyl-4H-1-benzopyran-2-propanoate and following the procedure of Example 9 gave methyl 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-4-oxo-8-propyl-4H-1-benzopyran-2-propanoate which was hydrolyzed using the procedure of Example 15 to give the title compound.

Microanalysis:

Calculated:

C, 68.62; H, 6.92.

Found:

C, 68.24; H, 6.92.

Example 23


5

Methyl 7-[[4-(4-acetyl-3-hydroxy-2-propylphenoxy)-2-butynyl]oxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoate

10

15

20



25

A solution of 2.04 g (7.8 mMole) of 4-(4-hydroxy-2-butynyloxy)-2-hydroxy-3-propylacetophenone, 2.49 g (7.8 mMole) of methyl (3,4-dihydro-7-hydroxy-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-yl)propanoate, and 2.04 g (7.8 mMole) of triphenylphosphine was prepared in 50 ml of dry tetrahydrofuran. After addition of 1.22 ml (7.8 mMole) of diethyl diazodicarboxylate the solution was stirred for 18 hours.

30

The tetrahydrofuran was removed by rotary evaporation and the residue stirred in 50 ml of ether. Insoluble solids were removed by filtration and the ether filtrate was concentrated by rotary evaporation to give 5.94 g of a crude oil. The oil was purified by elution chromatography to furnish 3.2 g (73%) of the methyl ester product.

Microanalysis:

Calculated:

C, 70.19; H, 7.14.

Found:

C, 70.33; H, 7.18

35

Example 24


40

Trans-methyl 7-[[4-(4-acetyl-3-hydroxy-2-propylphenoxy)-2-butenyl]oxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoate

45

50

55



A solution of 3.05 g (11.5 mMole) trans-4-(4-hydroxy-2-butenyloxy)-2-hydroxy-3-propylacetophenone, 3.70 g (11.5 mMole) of methyl 3-(3,4-dihydro-7-hydroxy-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-yl)-

propanoate, and 3.03 g (11.5 mMole) of triphenylphosphine in 75 ml of dry tetrahydrofuran was prepared. Diethyl diazodicarboxylate, 1.80 ml (11.5 mMole), was added and the solution was stirred for 18 hours at room temperature.

Solvent was removed from the reaction mixture by rotary evaporation and 75 ml of ether was added to the residue and stirred. After removal of insoluble solids by filtration, the ether filtrate was removed by rotary evaporation. The resulting crude material was purified by silica gel elution chromatography with 7% ethyl acetate/toluene to furnish 4.73 g (73%) of the methyl ester as an oil.

Microanalysis:

Calculated:


10 C, 69.94; H, 7.47.

Found:

C, 70.02; H, 7.45.

15 Example 25


20 Trans-7-[[4-(4-acetyl-3-methoxy-2-propylphenoxy)-2-butenyl]oxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoic acid



35 The compound of Example 24 is methylated using the method of Example 9 to give 7-[[4-(4-acetyl-3-methoxy-2-propylphenoxy)-2-butenyl]oxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoate which is then hydrolyzed using the method of Example 15 to give the title compound.

40 Example 26

Cis-methyl 7-[[4-(4-acetyl-3-hydroxy-2-propylphenoxy)-2-butenyl]oxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoate



15 A solution of 5.82 g (22 mMole) of cis-4-(4-hydroxy-2-butenyloxy)-2-hydroxy-3-propylacetophenone, 7.05 g (22 mMole) of methyl-(3,4-dihydro-7-hydroxy-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-yl)-propanoate, and 5.77 g (22 mMole) of triphenylphosphine was prepared in 150 ml of dry tetrahydrofuran. After addition of 3.5 ml (22 mMole) of diethyl diazodicarboxylate, the mixture was stirred for 18 hours.

20 The tetrahydrofuran was removed by rotary evaporation and the residue stirred with 100 ml of ether. Insoluble solids were removed by filtration and the ether filtrate was concentrated by rotary evaporation to give 16.6 g of yellow oil. The oil was purified by silica gel chromatography using 10% ethylacetate-toluene to furnish 6.28 g (50%) of pure methyl ester product.

Microanalysis:

Calculated:

25 C, 69.94; H, 7.47.

Found:

C, 69.52; H, 7.36.

30


Example 27

35

Cis-7-[[4-(4-acetyl-3-methoxy-2-propylphenoxy)-2-butenyl]oxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoic acid

40

45




50

The compound of Example 26 is methylated using the method of Example 9 to give cis-methyl 7-[[4-(4-acetyl-3-methoxy-2-propylphenoxy)-2-butenyl]-oxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoate which is then hydrolyzed using the method of Example 15 to give the title compound.

55

Example 28

(a) methyl 1,2,3,4-tetrahydro-7-methoxy-1-oxo-2-naphthalene carboxylate



28a

A 50% slurry of NaH (30g, 0.62 moles) in hexane was filtered through a fritted glass funnel to remove the mineral oil. The NaH was then added to a 2L flask, covered with 300ml of tetrahydrofuran (THF), and placed under a N₂ atmosphere. Upon the addition of 0.62 moles of dimethyl carbonate at one time with stirring, the reaction mixture was heated to 40°-50° C, whereupon 50g (0.28 moles) of commercially available 7-methoxy-1-tetralone in 150ml of THF was added at a rate to minimize foaming (1 hr.). After refluxing the reaction mixture for 2 hrs., the solution (red) was cooled to room temperature and slowly acidified by the addition of 45ml of acetic acid. The resulting paste was dissolved upon addition of 50ml of water. Ether was added and the layers were separated. The organic phase was washed with H₂O, 3% NaHCO₃ solution, and dried (Na₂SO₄). After filtration, the solvent was evaporated on a rotary evaporator. The residue was distilled at 168°-170° C at .2mm Hg.

Analysis for C₁₃H₁₆O₄ (MW = 236.26):


Calcd:

C, 66.66; H, 6.02.

Found:

C, 66.87; H, 6.07.

(b) methyl 1,2,3,4-tetrahydro-7-methoxy-2-naphthalenecarboxylate



28b

To 58g (.248 mole) of the product from Example 28(a) dissolved in a mixture of 387ml of acetic acid and 16.6ml of perchloric acid and placed in a 1L pressure bottle was added 5.8g of 5% Pd/C. The bottle was placed in a Parr shaker and hydrogenated at R.T. at 30 p.s.i. for 2 hr. After filtration, the filtrate was diluted with 1.1 liter of CHCl₃, washed with H₂O until the pH was neutral (5x). The organic phase was dried (Na₂SO₄) and the solvent evaporated to produce 49g of a crude yellow oil (the titled product).

Analysis for C₁₃H₁₆O₃ (MW = 220.26):


Calcd:

C, 70.88; H, 7.32.

Found:

C, 70.62; H, 7.09.

(c) 1,2,3,4-tetrahydro-7-methoxy-2-naphthalenecarboxylic acid



28c

To 49g (0.22 mole) of the product from Example 28(b) dissolved in 600ml of methanol at R.T. was added 150ml of 2M LiOH with stirring. After heating the stirred solution at 50 °C for 2 hr., the methanol was removed by rotary evaporation, the aqueous residue was washed 1x with ether, and acidified to pH 2 with HCl. Upon cooling and stirring, the crude acid precipitated and was separated by filtration. The crude acid was recrystallized from ether-hexane to give 30g (67%) of the product, m.p. 119° -122° C.

Analysis for $C_{12}H_{14}O_3$ (MW = 206.23):


Calcd:

C, 69.89; H, 6.84.

Found:

C, 69.70; H, 6.92.

(d) 1,2,3,4-tetrahydro-7-hydroxy-2-naphthalenecarboxylic acid



28d

To a 500ml flask containing 100g of pyridine hydrochloride was added 30g of the product of Example 28(c) and the reaction mixture was blanketed with N_2 . The reaction mixture was placed in an oil bath and heated to 215 °C for 2 hr. Upon cooling to R.T., 200ml of 1N HCl was added with stirring followed by 300ml of 1:1 mixture of ether/ethyl acetate. The organic phase was separated, washed 3x with H_2O , dried ($MgSO_4$), and filtered. Upon evaporation of the solvent, the residue was recrystallized from ether/hexane to produce 21.4g of the titled product, m.p. 170° -171° C.

Analysis for $C_{11}H_{12}O_3$ (MW = 192.21):


Calcd:

C, 68.73; H, 6.29.

Found:

C, 68.58; H, 6.35.

(e) methyl 1,2,3,4-tetrahydro-7-hydroxy-2-naphthalen carboxylate



28e

To 21g of the product of Example 28(d) dissolved in 250ml of methanol and placed in a 1L flask was added 30ml of trimethylorthoformate and 8ml of H_2SO_4 . The flask was covered and allowed to stand at room temperature for 3 days. Upon rotary evaporation of the methanol, 500ml of ether was added and the organic phase was washed 2x with 3% NaHCO_3 solution, and 2x with water. The organic phase was dried (Na_2SO_4), filtered, and the solvent removed by rotary evaporation. The residue was recrystallized from ethyl acetate-hexane to give 22g of the product, m.p. $83-84^\circ\text{C}$.

Analysis for $\text{C}_{12}\text{H}_{14}\text{O}_3$ (MW = 206.24):


Calcd:

C,69.89; H,6.84.

Found:

C,69.86; H,6.82.

(f) methyl 1,2,3,4-tetrahydro-7-(2-propenyloxy)-2-naphthalenecarboxylate



28f

To 20g (0.097 mole) of the product of Example 28(e) dissolved in 300ml of dry acetone was added 35g (2.5 eq.) of K_2CO_3 with vigorous stirring. An excess (20g) of allyl bromide was then added to the reaction vessel and the mixture was refluxed with a drying tube attached for 1 day. The reaction mixture was filtered and the solvent in the filtrate evaporated by rotary evaporation. The residue was taken up in 50ml of toluene and chromatographed on a 500g silica gel flash column, eluting with toluene. The eluent was evaporated and 17.5g (78%) of the product was isolated as a clear oil.

Analysis for $\text{C}_{15}\text{H}_{18}\text{O}_3$ (MW = 246.31):


Calcd:

C,73.15; H,7.37.

Found:

C,72.84; H,7.44.

(g) methyl 1,2,3,4-tetrahydro-7-hydroxy-8-(2-propenyl)-2-naphthalenecarboxylate



28g

To 20g of diethylaniline in a reaction vessel was added under argon 17g of the product of Example 28-
 (f). After the reaction mixture was heated at 215 °C for 6 hr., thin-layer chromatography (TLC) showed that
 two products had formed. Upon cooling, the reaction contents were poured into a 3:1 mixture of ethyl
 acetate/ether and the resulting solution washed with 200ml of 2N HCl and 100ml of water. The organic layer
 was separated, dried (Na₂SO₄), and filtered. Upon removal of the solvent by rotary evaporation, the crude
 residue (17g) was chromatographed on silica gel, eluting with 3% acetone/toluene. The most non-polar
 material (12.5g) was the titled product, 28g, mp. 90°-91° C. The most polar material was methyl 1,2,3,4-
 tetrahydro-7-hydroxy-6-(2-propenyl)-2-naphthalenecarboxylate.

Analysis for C₁₅H₁₈O₃ (MW = 246.31):


Calcd:

C, 73.15; H, 7.37.

Found:

C, 73.01; H, 7.34.

(h) methyl 1,2,3,4-tetrahydro-7-hydroxy-8-propyl-2-naphthalenecarboxylate



28h

To 12g of the product of Example 28(g) dissolved in 125ml of methanol and placed in a pressure bottle
 was added 1.25g of 5% Pd/C. The bottle was placed in a Parr shaker and the mixture hydrogenated at
 room temperature and 2 p.s.i. for 1 hr. Upon filtration of the reaction mixture and evaporation of the solvent,
 12g of the product 28h was produced as an oil.

Analysis for C₁₅H₂₀O₃ (MW = 248.32):

Calcd:


C, 72.55; H, 8.12.

Found:

C, 71.83; H, 8.20.

Example 29

7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxyl]-3,4-dihydro-4-hydroxy-8-propyl-2H-1-benzopyran-2-propanoic acid



Starting with methyl 7-[3-(4-acetyl-3-hydroxy-2-propylphenoxy)propoxyl]-3,4-dihydro-4-hydroxy-8-propyl-2H-1-benzopyran-2-propanoate prepared as disclosed in U. S. 4,665,203 and methylating the compound following the procedure of Example 9 then hydrolyzing that product according to the procedure of Example 15 gives the title compound.

Table 1 and Table 2 show test results for some embodiments of the invention. Table 3 shows the LTD₄ Receptor Binding test results.

Table 1

Compound	Receptor Binding		Chemotaxis ⁽²⁾	Degranulation
	LTB ₄	LTD ₄ ⁽¹⁾	(Boyden)	LTB ₄
	IC ₅₀	IC ₅₀	_____	IC ₅₀
Example 3	1.9x10 ⁻⁷ M	-	IC ₅₀ , 2x10 ⁻⁷ M	1.8x10 ⁻⁷ M
Example 4	5x10 ⁻⁶ M	-	62%(10 ⁻⁸ M)	2.7x10 ⁻⁶ M
Example 5	1x10 ⁻⁷ M	-	IC ₅₀ , 0.5x10 ⁻⁶ M	4x10 ⁻⁷ M
Example 6	2.6x10 ⁻⁶ M	Inactive	48%(10 ⁻⁶ M)	8x10 ⁻⁶ M
Example 7	0.5x10 ⁻⁶ M	Inactive	IC ₅₀ , 3.6x10 ⁻⁶ M	3.7x10 ⁻⁶ M
Example 8	1.8x10 ⁻⁶ M	-	12%(10 ⁻⁶ M)	-
Example 15	0.5x10 ⁻⁶ M	-	IC ₅₀ , 6.6x10 ⁻⁶ M	1.5x10 ⁻⁶ M
Example 16	0.25x10 ⁻⁶ M	Inactive	IC ₅₀ , 0.65x10 ⁻⁶ M	0.55x10 ⁻⁶ M
Example 17	0.8x10 ⁻⁶ M	-	31%(10 ⁻⁶ M)	2.1x10 ⁻⁶ M
Example 18	3.7x10 ⁻⁷ M	-	87%(10 ⁻⁵ M)	2.6x10 ⁻⁶ M
Example 19	1.2x10 ⁻⁶ M	-	15%(10 ⁻⁶ M)	5x10 ⁻⁶ M
Example 20	0.5x10 ⁻⁶ M	Inactive	IC ₅₀ , 1.7x10 ⁻⁶ M	0.4x10 ⁻⁷ M
Example 21	1x10 ⁻⁶ M	-	-	-
Example 1	0.48x10 ⁻⁶ M	Inactive	IC ₅₀ , 3x10 ⁻⁶ M	1.5x10 ⁻⁶ M
Example 22	0.5x10 ⁻⁶ M	-	IC ₅₀ , 1.2x10 ⁻⁶ M	1.6x10 ⁻⁶ M

(1) Inactive means the compound tested was inactive at 10⁻⁵M. These results are from Table 3.

(2) IC₅₀ is the effective concentration needed to cause 50% inhibition.

Table 2

<u>Compound</u>	<u>Guinea Pig Ileum Contraction pA₂</u>
Example 7	6.74
Example 15	Inactive
Example 20	6.5

Table 3LTD₄ Receptor Binding (1)LTB₄ Antagonist Testing Against an LTD₄ Receptor


<u>Compound [10 μM]</u>	<u>Percent Inhibition</u>		<u>Activity</u>
	<u>Test #1</u>	<u>Test #2</u>	
FPL 55712	54.68	46.96	A
Example 1	4.07	16.40	I
Example 20	+12.96	23.51	I
Example 16	16.49	24.08	I
Example 7	28.90	34.40	I
Example 6	+1.63	25.97	I

(1) This method employs LTD₄ membrane receptors prepared from a male guinea pig lung homogenate which was aliquoted and stored at -70°C. The particular preparation used to test the compounds of Table 3 was GPLP 9, having a K_D of 0.262 nM and yielding 0.447 pmol receptor/mg protein. Protein concentrations in these assays were adjusted to provide approximately 0.1 nM receptor in the presence of 1nM ³H-LTD₄. Incubation was conducted for 30 min. at 24°C and filtration methods were used to separate bound from


free ligand. Under these conditions, IC_{50} values for LTD_4 and FPL 55712, a known LTD_4 receptor antagonist, were $1.71 \times 10^{-8}M$ and $10.37 \times 10^{-6}M$ respectively. All compounds tested were initially solubilized in DMSO. Each compound was evaluated at a dose of $10 \mu M$ ($10^{-5}M$) in triplicate in 2 separate assays. The percent inhibition of LTD_4 specific binding for each compound is shown in Table 3. By the criteria established for potential LTD_4 receptor antagonists, compounds are active if the $10 \mu M$ test concentration inhibits LTD_4 specific binding by 45% or more. Using this parameter, all the Table 3 compounds were considered inactive relative to FPL 55712.

Scheme A and Scheme B show two preferred synthesis routes for making the compound of Formula III. Scheme C shows a general method for making compounds of the present invention.


EP 0 292 977 A1
SCHEME A



EP 0 292 977 A1
SCHEME B



SCHEME C




$y = 0$ to 6
 $x = 2$ to 7
 $B = CH_2$, $C = O$, $C-OH$
 $R^2 =$ methyl or ethyl
 $R^3 =$ alkyl
 $R^5 = H$, alkyl or
 R^5 and R^6 form a
 carbon to carbon bond.

$R^4 = H$ or alkyl
 $R^1 =$ alkyl, alkenyl, alkynyl, or
 $(CH_2)_n-R$ where $n = 1$ or 2
 and $R =$ cycloalkyl of 3-5
 carbon atoms.
 $R^8 =$ alkyl

Claims

1. A compound of the formula:



and the pharmaceutically acceptable salts thereof wherein

R¹ represents alkyl having 2-6 carbon atoms, alkenyl having 2 to 6 carbon atoms, alkynyl having 2 to 6 carbon atoms, or (CH₂)_nR wherein R represents cycloalkyl of 3 to 5 carbon atoms;

R² represents methyl or ethyl;

R³ represents alkyl having 1 to 5 carbon atoms;

W represents (CH₂)_x where x is 2 to 7, alkenylene having 3 to 7 carbon atoms, alkynylene having 3 to 7 carbon atoms, or cyclopentyl;

R⁴ represents hydrogen, alkyl having 2 to 5 carbon atoms, alkenyl having 2 to 5 carbon atoms, or alkynyl having 2 to 5 carbon atoms;

Q represents oxygen or CH₂;


B represents CH₂, C=O or CH-OH;

R⁵ represents hydrogen, alkyl having 1 to 6 carbon atoms, or R⁵ and R⁵ together optionally represent a carbon to carbon bond; or

R⁵ represents alkanoyl having 2 to 4 carbon atoms, carboxy, alkoxycarbonyl, or (CH₂)_y-CO₂R⁸ wherein y is 0 to 4 and R⁸ is hydrogen or alkyl having 1 to 6 carbon atoms; and

A represents -Z-CO₂R⁷ or -Z-CONR⁹R¹⁰ wherein R⁷ represents hydrogen or alkyl having 1 to 6 carbon atoms, R⁹ and R¹⁰ represent hydrogen, alkyl having 1 to 6 carbon atoms or cycloalkyl having 3 to 6 carbon atoms, or NR⁹R¹⁰ form a heterocyclic ring, and wherein Z is absent or represents straight or branched chain alkylene or alkenylene having up to 6 carbon atoms.

2. A compound according to Claim 1 of the formula



wherein

R¹ represents alkyl having 2-4 carbon atoms;

R² represents methyl or ethyl;

R³ represents alkyl having 1 to 3 carbon atoms;

W represents (CH₂)_x where x is 3 to 5, alkenylene having 3 to 5 carbon atoms, alkynylene having 3 to 5 carbon atoms, or cyclopentyl;


R⁴ represents alkyl having 2 to 4 carbon atoms;

R⁵ represents hydrogen, alkyl having 1 to 4 carbon atoms, or R⁵ and R⁵ together optionally represent a carbon to carbon bond;

A represents -Z-CO₂R⁷ or -Z-CONR⁹R¹⁰, wherein R⁷ represents hydrogen or alkyl having 1 to 4 carbon atoms, R⁹ and R¹⁰ represent hydrogen or alkyl having 1 to 4 carbon atoms, and wherein Z is absent or represents alkylene having up to 2 carbon atoms; and the stereoisomers and pharmaceutically

acceptable salts thereof.

3. A compound according to Claim 1 of the formula



wherein

R¹ represents alkyl having 2-4 carbon atoms;

R² represents methyl or ethyl;

R³ represents alkyl having 1 to 3 carbon atoms;


W represents (CH₂)_x where x is 3 to 5, alkenylene having 3 to 5 carbon atoms, alkynylene having 3 to 5 carbon atoms, or cyclopentyl;

R⁴ represents alkyl having 2 to 4 carbon atoms;

R⁵ represents hydrogen, alkyl having 1 to 4 carbon atoms, or R⁵ and R⁶ together optionally represent a carbon to carbon bond;


A represents -Z-CO₂R⁷ or -Z-CONR⁸R¹⁰, wherein R⁷ represents hydrogen or alkyl having 1 to 4 carbon atoms R⁸ and R¹⁰ represent hydrogen or alkyl having 1 to 4 carbons, and wherein Z is absent or represents alkylene having up to 2 carbon atoms; and the stereoisomers and pharmaceutically acceptable salts thereof.

4. A compound according to Claim 2 of the formula




wherein x is 3 to 5, R⁵ represents hydrogen or alkyl of 1 to 4 carbon atoms and A represents -Z-CO₂R⁷ wherein R⁷ represents hydrogen or alkyl having 1 to 4 carbon atoms and Z is absent or represents alkylene having up to 2 carbon atoms; and the stereoisomers and pharmaceutically acceptable salts thereof.

5. A compound according to Claim 4 of the formula




wherein p is 0 to 2; and the pharmaceutically acceptable salts thereof.

6. A compound according to Claim 3 of the formula




wherein x is 3 to 5, R⁵ represents hydrogen or alkyl of 1 to 4 carbon atoms or R⁵ and R⁶ together optionally represent a carbon to carbon bond; and A represents -Z-CO₂R⁷ wherein R⁷ represents hydrogen or alkyl having 1 to 4 carbon atoms and Z is absent or represents alkylene having up to 2 carbon atoms; and the stereoisomers and pharmaceutically acceptable salts thereof.

7. A compound according to Claim 6 of the formula



wherein R⁵ represents hydrogen or methyl or R⁵ and R⁶ together optionally represent a carbon to carbon bond; and wherein p is 0 to 2; and the pharmaceutically acceptable salts thereof.

8. A compound according to Claim 5 which is



and the pharmaceutically acceptable salts thereof.

9. A pharmaceutical composition comprising an effective anti-inflammatory amount of one of the compounds of Claims 1 to 8 in a pharmaceutical carrier.

10. An orally acceptable pharmaceutical composition of claim 9.

11. Use of one of the compounds of claims 1 to 8 for the production of a medicament against inflammatory diseases.

12. Use according to Claim 11 wherein the inflammatory diseases is rheumatoid arthritis, psoriasis, bowel disease or gout.

13. A compound according to Claim 2, which is 7-[3-[4-acetyl-2-(cyclopropylmethyl)-3-methoxyphenoxy]propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.

14. A compound according to Claim 2, which is 7-[[3-(4-acetyl-3-methoxy-2-propylphenoxy)-cyclopentyl]oxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.

15. A compound according to Claim 2, which is 7-[3-[4-acetyl-3-methoxy-2-(2-propenyl)phenoxy]propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.

16. A compound according to Claim 2, which is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid.

17. A compound according to Claim 3, which is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-4-oxo-8-propyl-4H-1-benzopyran-2-carboxylic acid.

18. A compound according to Claim 2, which is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)-propoxy]-3,4-dihydro-N-methyl-8-propyl-2H-1-benzopyran-2-carboxamide.

19. A compound according to Claim 2, which is methyl 7-[5-(4-acetyl-3-methoxy-2-propylphenoxy)-pentyloxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate.

20. A compound according to Claim 3, which is methyl 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)-propoxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoate.

21. A compound according to Claim 3, which is methyl 7-[[4-(4-acetyl-3-methoxy-2-propylphenoxy)-2-butynyl]oxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoate.

22. A compound according to Claim 1, which is methyl 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)-propoxy]-1,2,3,4-tetrahydro-8-propyl-2-naphthalenecarboxylate.

23. A compound according to Claim 2, which is methyl 3,4-dihydro-7-[3-[3-methoxy-4-(2-methyl-1-oxopropyl)-2-propylphenoxy]-propoxy]-8-propyl-2H-1-benzopyran-2-carboxylate.

24. A compound according to Claim 2, which is ethyl 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)-propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-yl-propanoate.

25. A compound according to Claim 2, which is 7-[5-(4-acetyl-3-methoxy-2-propylphenoxy)pentyloxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.

26. A compound according to Claim 3, which is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoic acid.

27. A compound according to Claim 3, which is 7-[[4-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-2-butynyl]oxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoic acid.

28. A compound according to Claim 2, which is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-1,2,3,4-tetrahydro-8-propyl-2-naphthalenecarboxylic acid.

29. A compound according to Claim 2, which is 3,4-dihydro-7-[3-[3-methoxy-4-(2-methyl-1-oxopropyl)-2-propylphenoxy]propoxy]-8-propyl-2H-1-benzopyran-2-carboxylic acid.

30. A compound according to Claim 2, which is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-propanoic acid.

31. A compound according to Claim 2, which is 7-[3-(4-acetyl-3-ethoxy-2-propylphenoxy)-propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.

32. A compound according to Claim 3, which is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-4-oxo-8-propyl-4H-1-benzopyran-2-propanoic acid.


33. A compound according to Claim 2, which is methyl 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)-propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate.

34. A pharmaceutical composition comprising an effective anti-inflammatory amount of one of the compounds of claims 13 to 33 in a pharmaceutical carrier.

35. Use of one of the compounds of Claims 13 to 33 for the production of a medicament against inflammatory diseases.

Claims for the following contracting State: GR

1. A compound of the formula:



and the pharmaceutically acceptable salts thereof wherein

R¹ represents alkyl having 2-6 carbon atoms, alkenyl having 2 to 6 carbon atoms, alkynyl having 2 to 6 carbon atoms, or (CH₂)_nR wherein R represents cycloalkyl of 3 to 5 carbon atoms;

R² represents methyl or ethyl;

R³ represents alkyl having 1 to 5 carbon atoms;

W represents (CH₂)_x where x is 2 to 7, alkenylene having 3 to 7 carbon atoms,

alkynylene having 3 to 7 carbon atoms, or cyclopentyl;

5 R⁴ represents hydrogen, alkyl having 2 to 5 carbon atoms, alkenyl having 2 to 5 carbon atoms, or alkynyl having 2 to 5 carbon atoms;


Q represents oxygen or CH₂;

B represents CH₂, C=O or CH-OH;

10 R⁵ represents hydrogen, alkyl having 1 to 6 carbon atoms, or R⁵ and R⁶ together optionally represent a carbon to carbon bond; or R⁵ represents alkanoyl having 2 to 4 carbon atoms, carboxy, alkoxy, carbonyl, or (CH₂)_y-CO₂R⁸ wherein y is 0 to 4 and R⁸ is hydrogen or alkyl having 1 to 6 carbon atoms; and

A represents -Z-CO₂R⁷ or -Z-CONR⁹R¹⁰ wherein R⁷ represents hydrogen or alkyl having 1 to 6 carbon atoms, R⁹ and R¹⁰ represent hydrogen, alkyl having 1 to 6 carbon atoms or cycloalkyl having 3 to 6 carbon atoms, or NR⁹R¹⁰ form a heterocyclic ring, and wherein Z is absent or represents straight or branched
15 chain alkylene or alkenylene having up to 6 carbon atoms.

2. A compound according to Claim 1 of the formula



wherein

30 R¹ represents alkyl having 2-4 carbon atoms;

R² represents methyl or ethyl;

R³ represents alkyl having 1 to 3 carbon atoms;


W represents (CH₂)_x where x is 3 to 5, alkenylene having 3 to 5 carbon atoms, alkynylene having 3 to 5 carbon atoms, or cyclopentyl;

35 R⁴ represents alkyl having 2 to 4 carbon atoms;

R⁵ represents hydrogen, alkyl having 1 to 4 carbon atoms, or R⁵ and R⁶ together optionally represent a carbon to carbon bond;

A represents -Z-CO₂R⁷ or -Z-CONR⁹R¹⁰, wherein R⁷ represents hydrogen or alkyl having 1 to 4 carbon atoms, R⁹ and R¹⁰ represent hydrogen or alkyl having 1 to 4 carbon atoms, and wherein Z is absent or represents alkylene having up to 2 carbon atoms; and the stereoisomers and pharmaceutically
40 acceptable salts thereof.

3. A compound according to Claim 1 of the formula



55 wherein

R¹ represents alkyl having 2-4 carbon atoms;

R² represents methyl or ethyl;

R³ represents alkyl having 1 to 3 carbon atoms;


W represents $(CH_2)_x$ where x is 3 to 5, alkenylene having 3 to 5 carbon atoms, alkynylene having 3 to 5 carbon atoms, or cyclopentyl;

R^4 represents alkyl having 2 to 4 carbon atoms;

R^5 represents hydrogen, alkyl having 1 to 4 carbon atoms, or R^5 and R^6 together optionally represent a carbon to carbon bond;


A represents $-Z-CO_2R^7$ or $-Z-CONR^8R^{10}$, wherein R^7 represents hydrogen or alkyl having 1 to 4 carbon atoms R^8 and R^{10} represent hydrogen or alkyl having 1 to 4 carbons, and wherein Z is absent or represents alkylene having up to 2 carbon atoms; and the stereoisomers and pharmaceutically acceptable salts thereof.

4. A compound according to Claim 2 of the formula




wherein x is 3 to 5, R^5 represents hydrogen or alkyl of 1 to 4 carbon atoms and A represents $-Z-CO_2R^7$ wherein R^7 represents hydrogen or alkyl having 1 to 4 carbon atoms and Z is absent or represents alkylene having up to 2 carbon atoms; and the stereoisomers and pharmaceutically acceptable salts thereof.

5. A compound according to Claim 4 of the formula




wherein p is 0 to 2; and the pharmaceutically acceptable salts thereof.

6. A compound according to Claim 3 of the formula




wherein x is 3 to 5, R^5 represents hydrogen or alkyl of 1 to 4 carbon atoms or R^5 and R^6 together optionally represent a carbon to carbon bond; and A represents $-Z-CO_2R^7$ wherein R^7 represents hydrogen or alkyl having 1 to 4 carbon atoms and Z is absent or represents alkylene having up to 2 carbon atoms; and the stereoisomers and pharmaceutically acceptable salts thereof.

7. A compound according to Claim 6 of the formula



wherein R^5 represents hydrogen or methyl or R^5 and R^6 together optionally represent a carbon to carbon bond; and wherein p is 0 to 2; and the pharmaceutically acceptable salts thereof.

8. A compound according to Claim 5 which is



and the pharmaceutically acceptable salts thereof.

9. A compound according to Claim 2, which is 7-[3-(4-acetyl-2-(cyclopropylmethyl)-3-methoxyphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.

10. A compound according to Claim 2, which is 7-[[3-(4-acetyl-3-methoxy-2-propylphenoxy)-cyclopentyl]oxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.

11. A compound according to Claim 2, which is 7-[3-(4-acetyl-3-methoxy-2-(2-propenyl)phenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.

12. A compound according to Claim 2, which is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid.

13. A compound according to Claim 3, which is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-4-oxo-8-propyl-4H-1-benzopyran-2-carboxylic acid.

14. A compound according to Claim 2, which is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-N-methyl-8-propyl-2H-1-benzopyran-2-carboxamide.

15. A compound according to Claim 2, which is methyl 7-[5-(4-acetyl-3-methoxy-2-propylphenoxy)pentyl]oxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate.

16. A compound according to Claim 3, which is methyl 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoate.

17. A compound according to Claim 3, which is methyl 7-[[4-(4-acetyl-3-methoxy-2-propylphenoxy)-2-butynyl]oxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoate.

18. A compound according to Claim 1, which is methyl 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-1,2,3,4-tetrahydro-8-propyl-2-naphthalenecarboxylate.

19. A compound according to Claim 2, which is methyl 3,4-dihydro-7-[8-[3-methoxy-4-(2-methyl-1-oxopropyl)-2-propylphenoxy]propoxy]-8-propyl-2H-1-benzopyran-2-carboxylate.

20. A compound according to Claim 2, which is ethyl 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-yl-propanoate.

21. A compound according to Claim 2, which is 7-[[5-(4-acetyl-3-methoxy-2-propylphenoxy)pentyl]oxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.

22. A compound according to Claim 3, which is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoic acid.

23. A compound according to Claim 3, which is 7-[[4-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-2-butynyl]oxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoic acid.

24. A compound according to Claim 2, which is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-1,2,3,4-tetrahydro-8-propyl-2-naphthalenecarboxylic acid.

25. A compound according to Claim 2, which is 3,4-dihydro-7-[3-[3-methoxy-4-(2-methyl-1-oxopropyl)-2-propylphenoxy]propoxy]-8-propyl-2H-1-benzopyran-2-carboxylic acid.

26. A compound according to Claim 2, which is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-propanoic acid.


27. A compound according to Claim 2, which is 7-[3-(4-acetyl-3-ethoxy-2-propylphenoxy)-propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.

28. A compound according to Claim 3, which is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-4-oxo-8-propyl-4H-1-benzopyran-2-propanoic acid.

29. A compound according to Claim 2, which is methyl 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)-propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate.

15 Claims for the following contracting State: ES

1. A process for the production of a compound of the formula:



and the pharmaceutically acceptable salts thereof wherein

R¹ represents alkyl having 2-6 carbon atoms, alkenyl having 2 to 6 carbon atoms, alkynyl having 2 to 6 carbon atoms, or (CH₂)_nR wherein R represents cycloalkyl of 3 to 5 carbon atoms;

R² represents methyl or ethyl;

R³ represents alkyl having 1 to 5 carbon atoms;

W represents (CH₂)_x where x is 2 to 7, alkenylene having 3 to 7 carbon atoms, alkynylene having 3 to 7 carbon atoms, or cyclopentyl;


R⁴ represents hydrogen, alkyl having 2 to 5 carbon atoms, alkenyl having 2 to 5 carbon atoms, or alkynyl having 2 to 5 carbon atoms;

Q represents oxygen or CH₂;

B represents CH₂, C=O or CH-OH;

R⁵ represents hydrogen, alkyl having 1 to 6 carbon atoms, or R⁵ and R⁶ together optionally represent a carbon to carbon bond; or R⁵ represents alkanoyl having 2 to 4 carbon atoms, carboxyl, alkoxycarbonyl, or (CH₂)_y-CO₂R⁸ wherein y is 0 to 4 and R⁸ is hydrogen or alkyl having 1 to 6 carbon atoms; and

A represents -Z-CO₂R⁷ or -Z-CONR⁹R¹⁰ wherein R⁷ represents hydrogen or alkyl having 1 to 6 carbon atoms, R⁹ and R¹⁰ represent hydrogen, alkyl having 1 to 6 carbon atoms or cycloalkyl having 3 to 6 carbon atoms, or NR⁹R¹⁰ form a heterocyclic ring, and wherein Z is absent or represents straight or branched chain alkylene or alkenylene having up to 6 carbon atoms, characterized in that a compound of the formula




wherein R_1 , R_3 , W, R_4 , Q, B, R_5 , A and R_6 are defined as above, is alkylated; or a phenone intermediate is alkylated prior to forming the -O-W-O-bridge.

2. A process according to Claim 1 wherein the produced compound has the formula

5

10



15 wherein

R^1 represents alkyl having 2-4 carbon atoms;

R^2 represents methyl or ethyl;

R^3 represents alkyl having 1 to 3 carbon atoms;

W represents $(CH_2)_x$ where x is 3 to 5, alkenylene having 3 to 5 carbon atoms, alkynylene having 3 to 5 carbon atoms, or cyclopentyl;

R^4 represents alkyl having 2 to 4 carbon atoms;


R^5 represents hydrogen, alkyl having 1 to 4 carbon atoms, or R^5 and R^6 together optionally represent a carbon to carbon bond;

A represents $-Z-CO_2R^7$ or $-Z-CONR^9R^{10}$, wherein R^7 represents hydrogen or alkyl having 1 to 4 carbon atoms, R^9 and R^{10} represent hydrogen or alkyl having 1 to 4 carbon atoms, and wherein Z is absent or represents alkylene having up to 2 carbon atoms; and the stereoisomers and pharmaceutically acceptable salts thereof.

3. A process according to Claim 1 wherein the produced compound has the formula

30

35



40

45 wherein

R^1 represents alkyl having 2-4 carbon atoms;

R^2 represents methyl or ethyl;

R^3 represents alkyl having 1 to 3 carbon atoms;

W represents $(CH_2)_x$ where x is 3 to 5, alkenylene having 3 to 5 carbon atoms, alkynylene having 3 to 5 carbon atoms, or cyclopentyl;

R^4 represents alkyl having 2 to 4 carbon atoms;


R^5 represents hydrogen, alkyl having 1 to 4 carbon atoms, or R^5 and R^6 together optionally represent a carbon to carbon bond;

A represents $-Z-CO_2R^7$ or $-Z-CONR^9R^{10}$, wherein R^7 represents hydrogen or alkyl having 1 to 4 carbon atoms, R^9 and R^{10} represent hydrogen or alkyl having 1 to 4 carbons, and wherein Z is absent or represents alkylene having up to 2 carbon atoms; and the stereoisomers and pharmaceutically acceptable salts thereof.

50


4. A process according to Claim 2 wherein the produced compound has the formula

55




10 wherein x is 3 to 5, R⁵ represents hydrogen or alkyl of 1 to 4 carbon atoms and A represents -Z-CO₂R⁷ wherein R⁷ represents hydrogen or alkyl having 1 to 4 carbon atoms and Z is absent or represents alkylene having up to 2 carbon atoms; and the stereoisomers and pharmaceutically acceptable salts thereof.

5. A process according to Claim 4 wherein the produced compound has the formula




25 wherein p is 0 to 2; and the pharmaceutically acceptable salts thereof.

6. A process according to Claim 3 wherein the produced compound has the formula




40 wherein x is 3 to 5, R⁵ represents hydrogen or alkyl of 1 to 4 carbon atoms or R⁵ and R⁶ together optionally represent a carbon to carbon bond; and A represents -Z-CO₂R⁷ wherein R⁷ represents hydrogen or alkyl having 1 to 4 carbon atoms and Z is absent or represents alkylene having up to 2 carbon atoms; and the stereoisomers and pharmaceutically acceptable salts thereof.

7. A process according to Claim 6 wherein the produced compound has the formula



55 wherein R⁵ represents hydrogen or methyl or R⁵ and R⁶ together optionally represent a carbon to carbon bond; and wherein p is 0 to 2; and the pharmaceutically acceptable salts thereof.

8. A process according to Claim 5 wherein the produced compound is



and the pharmaceutically acceptable salts thereof.

9. A process according to Claim 2 wherein the produced compound is 7-[3-(4-acetyl-2-(cyclopropylmethyl)-3-methoxyphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.

10. A process according to Claim 2 wherein the produced compound is 7-[[3-(4-acetyl-3-methoxy-2-propylphenoxy)cyclopentyl]oxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.

11. A process according to Claim 2 wherein the produced compound is 7-[3-(4-acetyl-3-methoxy-2-(2-propenyl)phenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.

12. A process according to Claim 2 wherein the produced compound is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-2H-1-benzopyran-2-carboxylic acid.

13. A process according to Claim 3 wherein the produced compound is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-4-oxo-8-propyl-4H-1-benzopyran-2-carboxylic acid.

14. A process according to Claim 2 wherein the produced compound is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-N-methyl-8-propyl-2H-1-benzopyran-2-carboxamide.

15. A process according to Claim 2 wherein the produced compound is methyl 7-[5-(4-acetyl-3-methoxy-2-propylphenoxy)pentyl]oxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate.

16. A process according to Claim 3 wherein the produced compound is methyl 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoate.

17. A process according to Claim 3 wherein the produced compound is methyl 7-[[4-(4-acetyl-3-methoxy-2-propylphenoxy)-2-butynyl]oxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoate.

18. A process according to Claim 1 wherein the produced compound is methyl 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-1,2,3,4-tetrahydro-8-propyl-2-naphthalenecarboxylate.

19. A process according to Claim 2 wherein the produced compound is methyl 3,4-dihydro-7-[3-[3-methoxy-4-(2-methyl-1-oxopropyl)-2-propylphenoxy]propoxy]-8-propyl-2H-1-benzopyran-2-carboxylate.

20. A process according to Claim 2 wherein the produced compound is ethyl 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-yl-propanoate.

21. A process according to Claim 2 wherein the produced compound is 7-[[5-(4-acetyl-3-methoxy-2-propylphenoxy)pentyl]oxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.

22. A process according to Claim 3 wherein the produced compound is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoic acid.

23. A process according to Claim 3 wherein the produced compound is 7-[[4-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-2-butynyl]oxy]-3,4-dihydro-2-methyl-4-oxo-8-propyl-2H-1-benzopyran-2-propanoic acid.

24. A process according to Claim 2 wherein the produced compound is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-1,2,3,4-tetrahydro-8-propyl-2-naphthalenecarboxylic acid.

25. A process according to Claim 2 wherein the produced compound is 3,4-dihydro-7-[3-[3-methoxy-4-(2-methyl-1-oxopropyl)-2-propylphenoxy]propoxy]-8-propyl-2H-1-benzopyran-2-carboxylic acid.

26. A process according to Claim 2 wherein the produced compound is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-propanoic acid.

27. A process according to Claim 2 wherein the produced compound is 7-[3-(4-acetyl-3-ethoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylic acid.

28. A process according to Claim 3 wherein the produced compound is 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-4-oxo-8-propyl-4H-1-benzopyran-2-propanoic acid.

29. A process according to Claim 2 wherein the produced compound is methyl 7-[3-(4-acetyl-3-methoxy-2-propylphenoxy)propoxy]-3,4-dihydro-8-propyl-2H-1-benzopyran-2-carboxylate.



European Patent
Office

EUROPEAN SEARCH REPORT

Application number

DOCUMENTS CONSIDERED TO BE RELEVANT			EP 88108449.5
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
A	EP - B1 - 0 017 332 (FISONS) * Claims 1,12,13 * --	1,9,11	C 07 D 311/22 C 07 D 311/24 C 07 D 311/58
A	US - A - 4 546 194 (MIYANO et al.) * Claim 1; column 15, lines 37-44 * --	1,9,11	C 07 D 311/66 A 61 K 31/35 C 07 C 57/48 C 07 C 61/39
A	CHEMICAL ABSTRACTS, vol. 103, no. 19, November 11, 1985, Columbus, Ohio, USA SEARLE, G.D. "Benzopyran antimetabolites" page 699, column 1, abstract-no. 160 389g & Jpn.Kokai Tokkyo Koho JP 60-42 378 [85-42 378] --	1,9,11	
A	CHEMICAL ABSTRACTS, vol. 97, no. 3, July 19, 1982, Columbus, Ohio, USA SHEARD, P.; HOLROYDE, M.C.; GHELANI, A.M.; BANTICK, J.R.; LEE, T.B. "Antagonists of SRS-A and leukotrienes" page 44, column 1, abstract-no. 16 807g & Adv. Prostaglandin, Thromboxane, Leukotriene Res. 1982, 9(Leukotrienes Other Lipxygenase Prod.), 229-35 ----	1,9,11	TECHNICAL FIELDS SEARCHED (Int. Cl.4) C 07 D 311/00 C 07 C 57/00 C 07 C 61/00
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 09-09-1988	Examiner BRUS
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	